



National Library
of Canada

Bibliothèque nationale
du Canada

Canadian Theses Service

Service des thèses canadiennes

Ottawa, Canada
K1A 0N4

NOTICE

The quality of this microform is heavily dependent upon the quality of the original thesis submitted for microfilming. Every effort has been made to ensure the highest quality of reproduction possible.

If pages are missing, contact the university which granted the degree.

Some pages may have indistinct print especially if the original pages were typed with a poor typewriter ribbon or if the university sent us an inferior photocopy.

Reproduction in full or in part of this microform is governed by the Canadian Copyright Act, R.S.C. 1970, c. C-30, and subsequent amendments.

AVIS

La qualité de cette microforme dépend grandement de la qualité de la thèse soumise au microfilmage. Nous avons tout fait pour assurer une qualité supérieure de reproduction.

S'il manque des pages, veuillez communiquer avec l'université qui a conféré le grade.

La qualité d'impression de certaines pages peut laisser à désirer, surtout si les pages originales ont été dactylographiées à l'aide d'un ruban usé ou si l'université nous a fait parvenir une photocopie de qualité inférieure.

La reproduction, même partielle, de cette microforme est soumise à la Loi canadienne sur le droit d'auteur, SRC 1970, c. C-30, et ses amendements subséquents.

UNIVERSITY OF ALBERTA
THE QUANTUM WITT ALGEBRA
AND
QUANTIZATIONS OF SOME MODULES OVER THE WITT ALGEBRA

by



KE-QIN LIU

A THESIS
SUBMITTED TO THE FACULTY OF GRADUATE STUDIES AND RESEARCH
IN PARTIAL FULFILLMENT OF THE REQUIREMENTS
FOR THE DEGREE OF
DOCTOR OF PHILOSOPHY

DEPARTMENT OF MATHEMATICS

EDMONTON, ALBERTA
SPRING, 1992



National Library
of Canada

Bibliothèque nationale
du Canada

Canadian Theses Service Service des thèses canadiennes

Ottawa, Canada
K1A 0N4

The author has granted an irrevocable non-exclusive licence allowing the National Library of Canada to reproduce, loan, distribute or sell copies of his/her thesis by any means and in any form or format, making this thesis available to interested persons.

The author retains ownership of the copyright in his/her thesis. Neither the thesis nor substantial extracts from it may be printed or otherwise reproduced without his/her permission.

L'auteur a accordé une licence irrévocable et non exclusive permettant à la Bibliothèque nationale du Canada de reproduire, prêter, distribuer ou vendre des copies de sa thèse de quelque manière et sous quelque forme que ce soit pour mettre des exemplaires de cette thèse à la disposition des personnes intéressées.

L'auteur conserve la propriété du droit d'auteur qui protège sa thèse. Ni la thèse ni des extraits substantiels de celle-ci ne doivent être imprimés ou autrement reproduits sans son autorisation.

ISBN 0-315-73103-6

Canada

UNIVERSITY OF ALBERTA

RELEASE FORM

NAME OF AUTHOR: KE-QIN LIU


TITLE OF THESIS: THE QUANTUM WITT ALGEBRA AND QUANTIZATION OF SOME
MODULES OVER THE WITT ALGEBRA

DEGREE FOR WHICH THESIS WAS PRESENTED: DOCTOR OF PHILOSOPHY

YEAR THIS DEGREE GRANTED 1992

Permission is hereby granted to the UNIVERSITY OF ALBERTA LIBRARY to reproduce single copies of the thesis and to lend or sell such copies for private, scholarly or scientific research purposes only.

The author reserves other publication rights, and neither the thesis nor extensive extracts from it may be printed or otherwise reproduced without the author's written permission.



SIGNED

PERMANENT ADDRESS:

Department of Mathematics

University of Alberta


Edmonton, Canada, T6G 2G1

Date



Jan. 24, 1992

UNIVERSITY OF ALBERTA
THE FACULTY OF GRADUATE STUDIES AND RESEARCH


The undersigned certify that they have read, and recommend to the Faculty of Graduate Studies and Research, for acceptance, a thesis entitled THE QUANTUM WITT ALGEBRA AND QUANTIZATIONS OF SOME MODULES OVER THE WITT ALGEBRA submitted by KE-QIN LIU in partial fulfillment of the degree of Doctor of Philosophy.




R.V. Moody
Supervisor


A. Pianzola

A.N. Al-Hussaini




T.D. Rogers



M. Shirvani

Date

Dec. 12, 1991


K. Misra

ABSTRACT

In this thesis, we investigate the problem of quantizing the Witt algebra and its representations over complex number field.

In chapter 1, we begin by introducing the quantum Witt algebra, the quantum Virasoro algebra and a q -analogue of the simplest affine Kac-Moody algebra $sl(2)$. Each of these objects is an example of a new class of non-associative algebras. We call this class of non-associative algebras the quantum Lie algebras, which are defined by using a q -analogue of the Jacobi identity. Next, we introduce the concept of the quantum universal central extension of a quantum Lie algebra and study its basic properties. Finally, we determine the quantum universal central extension of some quantum Lie algebras.

In chapter 2, after introducing quantum flexible algebras, we prove that q -analogues of the two characterizations of the usual Witt algebra hold for the quantum Witt algebra.

In chapter 3, we give our quantization of the enveloping algebra of the Witt algebra; construct the q -analogues of the module of tensor fields over the Witt algebra; study their properties and prove a q -analogue of Kaplansky's theorem concerning the module of tensor fields over the Witt algebra.

ACKNOWLEDGEMENT

I would like to take this opportunity to thank my supervisor Prof. Robert V. Moody firstly for accepting me as his student and secondly for his guidance, encouragement and interest in my work.

I am also grateful to Prof. Kailash C. Misra, Prof. Arturo Pianzola and Dr. Maziar Shirvani for their helpful comments.

TABLE OF CONTENTS

Chapter 0. Introduction	1
References	4
Chapter 1. Quantum central extensions	5
§1.0 Introduction	5
§1.1 J -derivations and the quantum Witt algebra	5
§1.2 Quantum central extensions	11
§1.3 The quantum universal central extension of W_q and $C[t, t^{-1}] \otimes s\ell_q^n(2)$	19
References	24
Chapter 2. The characterizations of the quantum Witt algebra	25
§2.0 Basic definitions and main threorem	25
§2.1 The proof of the theorem $:(2) \Rightarrow (1)$	27
§2.2 The proof of the theorem $:(3) \Rightarrow (1)$	31
References	35
Chapter 3. Quantizations of the module of tensor fields over the Witt algebra	36
§3.0 Introduction	36
§3.1 The constructions of $U(W_q)$ -modules $X(\lambda, \alpha, \beta)$	36
§3.2 The properties of $U(W_q)$ -modules $X(\lambda, \alpha, \beta)$	40
§3.3 A q -analogue of Kaplansky's theorem	54
References	62
Conclusions and some open problems	63
References	64

CHAPTER 0. INTRODUCTION

The search for q -analogues of known mathematical objects, which is affectionately called the " q -disease", has a long history in mathematics. As early as the 1840's, A.-L. Cauchy, E. Heine and other mathematicians found a q -analogue of the binomial theorem for $|q| < 1$. In 1869, J. Thomae introduced a q -analogue of the gamma function for $0 < q < 1$. The 19th century mathematician F. H. Jackson devoted much of his mathematical career to developing q -analogues of the hypergeometric series. Over the past fifteen years, R. Askey, his students and his collaborators have produced a substantial amount of interesting work in the theory of q -hypergeometric series. However, the general area of algebra was not infected by the q -disease until just ten years ago.

In 1981, P. Kulish and N. Reshetikhin [11] introduced the C -associative algebra $U_h(sl(2))$ with the generators X^+ , X^- , H and the relations:

$$\begin{aligned}HX^\pm - X^\pm H &= \pm 2X^\pm, \\ X^+X^- - X^-X^+ &= \frac{e^{hH} - e^{-hH}}{e^h - e^{-h}}, \quad h \in \mathbb{C},\end{aligned}$$

which arose from their study of Quantum Inverse Scattering Method. The associative algebra $U_h(sl(2))$ can be regarded as a q -analogue of the enveloping algebra of the 3-dimensional simple Lie algebra $sl(2)$ because when $q := e^h$ approaches unity, $U_h(sl(2))$ becomes the enveloping algebra of $sl(2)$. It was this object that V. Drinfeld [2,3] and M. Jimbo [6,7] recently generalized to the case of finite dimensional simple Lie algebras.

The main development in the representation theory of Lie algebras have been with the finite dimensional Lie algebras, the Kac-Moody algebras and the Virasoro algebra. Although the enveloping algebras of the finite dimensional simple Lie algebras and the Kac-Moody algebras have been quantized and q -analogues of their representations have been found, this is not case for the Virasoro algebra.

It is well-known that the representations of the Virasoro algebra with the center acting trivially, i.e., the representations of the Witt algebra, play a fundamental role in the representation theory of the Virasoro algebra. Therefore, the first problem in developing q -analogues of the representations of the Virasoro algebra is naturally to find q -analogues of the representations of the Witt algebra. The purpose of my thesis is to first address the problem of quantizing the Witt algebra, the Virasoro algebra and their enveloping algebras, and then to investigate possible q -analogues of the representations of the Witt algebra.

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$ -TeX

What follows is a summary of the results of my thesis.

Let q be a fixed non-zero complex number with $q^2 \neq 1$ and let $[m]$ denote the " q -integer" $\frac{q^m - q^{-m}}{q - q^{-1}}$, where $m \in \mathbb{Z}$. Our quantization of the Witt algebra and its enveloping algebra originated by considering certain operators, $\{D_n \mid n \in \mathbb{Z}\}$, called J -derivations on the Laurent polynomial ring $\mathbb{C}[t, t^{-1}]$, where the operators J and $\{D_n \mid n \in \mathbb{Z}\}$ defined by

$$\begin{aligned} J : t^n &\mapsto q^n t^n, \\ \left(\frac{d}{dt}\right)_q : t^n &\mapsto [n] t^{n-1}, \\ D_n &:= -t^{n+1} \left(\frac{d}{dt}\right)_q. \end{aligned}$$

It turns out that $\mathcal{D} := \oplus_{n \in \mathbb{Z}} \mathbb{C} D_n$ consists of all J -derivations of $\mathbb{C}[t, t^{-1}]$ and the bracket on $\mathcal{D}$, which is defined by

$$(0.1) \quad [D_m, D_n] := J D_m J^{-1} D_n J - J D_n J^{-1} D_m J \quad \text{for } m, n \in \mathbb{Z},$$

satisfies

$$[D_m, D_n] = [m - n] D_{m+n}.$$

Clearly, these bracket relations can be obtained from those of the usual Witt algebra by replacing the structure constants $\{m - n \mid m, n \in \mathbb{Z}\}$ by their q -analogues $\{[m - n] \mid m, n \in \mathbb{Z}\}$. So it seems reasonable to take $\mathcal{D}$, with the bracket (0.1), as a realization of the q -analogue of the usual Witt algebra. This produces the **quantum Witt algebra**.

Parallel to the fact that the Lie bracket on the Witt algebra satisfies the Jacobi identity, we will prove that the bracket on $\mathcal{D}$ above satisfies the following q -analogue of the Jacobi identity:

$$(0.2) \quad [[x, y], \sigma(z)] + [[y, z], \sigma(x)] + [[z, x], \sigma(y)] = 0 \quad \text{for } x, y, z \in \mathcal{D},$$

where $\sigma : \mathcal{D} \rightarrow \mathcal{D}$ is a linear map defined by

$$\sigma(D_m) := \frac{q^m + q^{-m}}{2} D_m \quad \text{for } m \in \mathbb{Z}.$$

We call (0.2) the **quantum Jacobi identity**.

Using the quantum Jacobi identity, we will introduce a class of non-associative algebras called **quantum Lie algebras**, which are a generalization of Lie algebras. In order to discover the q -analogue of the Virasoro algebra, we

will define the **quantum universal central extensions** (q -U.C.E.) of quantum Lie algebras and study the q -U.C.E. of the quantum Witt algebra. We will find that if q is not a root of unity, then the q -U.C.E. of the quantum Witt algebra is a 1-dimensional quantum central extension, and this object is defined to be the **quantum Virasoro algebra**.

This completes our summary of the main development in chapter 1.

Before describing the contents of chapter 2, let us recall some interesting results about the Witt algebra. From H.C. Myung's book [14] on Malcev-admissible algebras and I. Kaplansky's paper [9] on the Virasoro algebra, we know that if $A = \bigoplus_{n \in \mathbb{Z}} A_n$ is a $\mathbb{Z}$ -graded algebra over complex number field, then the following are equivalent:

- A is the Witt algebra.
- A is a flexible algebra and the attached minus algebra $(A^-, [-, -])$ is the Witt algebra.
- A is a Lie algebra, $\dim(A_n) \leq 1$, $n \in \mathbb{Z}$ and

$$(A_0 A_1) \neq 0, (A_1 A_{-1}) \neq 0, (A_2 A_{-1}) \neq 0, (A_1 A_{-2}) \neq 0.$$

In chapter 2, after introducing the concept of the **quantum flexible algebras**, we will prove that if q is not a root of unity, then the quantizations of the results above hold for the quantum Witt algebra. It is worth pointing out that throughout this work, the quantum Jacobi identity plays an essential role and we feel that the quantum Lie algebras are a class of non-associative algebras which deserve more attention.

In chapter 3, we will investigate possible q -analogues of the representations of the Witt algebra. The most important representation of the Witt algebra is the one called the module of tensor fields $V_{\alpha\beta}$, where α and β are complex numbers. In 1982, B.L. Feigin and D.B. Fuchs [4] introduced this module and used it in their proof of the Kac's determinant formula. In the same year, I. Kaplansky [9] proved the following remarkable property of the module of tensor fields $V_{\alpha\beta}$:

- If $V = \bigoplus_{n \in \mathbb{Z}} C v_n$ is a $\mathbb{Z}$ -graded module over the Witt algebra $W = \bigoplus_{n \in \mathbb{Z}} C d_n$ and $d_{\pm 1}$ are injective operators on V , then V is isomorphic to the module of tensor fields $V_{\alpha\beta}$ for some complex numbers α and β .

We call this result Kaplansky's theorem. The basic idea in the proof of Kaplansky's theorem has greatly influenced mathematicians and physicists who attacked Kac's conjecture concerning the representations of the Virasoro algebra (see [1], [10] and [12]). Therefore, we consider quantizing Kaplansky's theorem an important first step in developing q -analogues of the representations of the Virasoro algebra.

Motivated by our work in chapter 1, we will define the q -analogue of the universal enveloping algebra of the Witt algebra, denoted by $U(W_q)$. Using the

operations on the $\mathbb{Z}$ -graded modules over the Witt algebra in [4], we will construct two kinds of $U(W_q)$ -modules $A(\lambda, \alpha, \beta)$ and $B(\lambda, \alpha, \beta)$, where $\lambda, \alpha, \beta \in \mathbb{C}$ and $\lambda \neq 0$. Both $A(\lambda, \alpha, \beta)$ and $B(\lambda, \alpha, \beta)$ are q -analogues of the module of tensor fields. It turns out that under conditions similar to the ones in Kaplansky's theorem, every $\mathbb{Z}$ -graded $U(W_q)$ -module $V = \bigoplus_{n \in \mathbb{Z}} C v_n$ is isomorphic to either $A(\lambda, \alpha, \beta)$ or $B(\lambda, \alpha, \beta)$ for some $\lambda, \alpha, \beta \in \mathbb{C}$ and $\lambda \neq 0$. This is our quantization of Kaplansky's theorem obtained in chapter 3.

REFERENCES

1. V.Chari and A.Pressley, *Unitary representations of the Virasoro algebra and a conjecture of Kac*, *Compositio Mathematica* **67** (1989), 315-342.
2. V.Drinfeld, *Dedekind ANSHEB* **243**, **NS** (1985), 1001-1004.
3. V.Drinfeld, *Proc. of the ICM-86 Berkeley, v.1* (1986), 738-739.
4. B.L.Figini and D.B.Fuchs, *Invariant skew-symmetric differential operators on the line and Verma modules over the Virasoro algebra*, *Funct. Anal. Appl.* (English translation) **16** (1982), 114-126.
5. G.Gasper and M.Rahman, "Basic hypergeometric series," Cambridge university press, 1990.
6. M.Jimbo, *Commun. Math. Phys.* **102** (1986), 535-547.
7. M.Jimbo, *Lett. Math. Phys.* **11** (1986), 247-252.
8. V.G.Kac, "Infinite dimensional Lie algebras," Cambridge university press.
9. I.Kaplansky, *The Virasoro algebra*, *Commun. Math. Phys.* **86** (1982), 49-54.
10. I.Kaplansky and L.J.Santharoubane, *Harish-Chandra modules over the Virasoro algebra*, *Proc. Conf. "Infinite dimensional Lie groups"* . MSRI Publications (1985), 217-231.
11. P.Kulish and N.Reshetikhin, *Zap. nauchn. seminarov LOMI* **101** (1981), 101-110.
12. C.Martin and A.Piard, *Indecomposable modules over the Virasoro Lie algebra and a conjecture of V.Kac*, *Commun. Math. Phys.* **137** (1991), 109-132.
13. R.V.Moody and A.Pianzola, "Lie algebras with triangular decomposition," preprint.
14. H.C.Myung, "Malcev-admissible algebras," *Progress in mathematics*, Birkhäuser Boston, Inc., 1986.
15. L.A.Takhtajan, *Introduction to quantum groups*, *Lecture Note in Physics* (370), 3-28.

CHAPTER 1. QUANTUM CENTRAL EXTENSIONS

§1.0. Introduction

The purpose of this chapter is to give one possible solution to some questions proposed by Y.I. Manin in [6]. In section 1, we will construct the quantum Witt algebra W_q from the operators called J -derivations on the Laurent polynomial ring $C[t, t^{-1}]$; it turns out that W_q satisfies the quantum Jacobi identity. Other objects which satisfy the quantum Jacobi identity are the quantum Virasoro algebra V_q (where q is not a root of unity and $q \neq 0$) and the quantum Kac-Moody algebra $\widehat{\mathfrak{sl}}_q^n(2)$ (where n is a positive integer and $q^{2n} \neq 0, 1$). All these examples are q -analogues of familiar objects and become those objects when $q \rightarrow 1$. In section 2, we will define quantum universal central extension and study its basic properties. In section 3, we will prove that quantum Virasoro algebra V_q and the quantum Kac-Moody algebra $\widehat{\mathfrak{sl}}_q^n(2)$ are the quantum universal central extensions of W_q and $C[t, t^{-1}] \otimes \widehat{\mathfrak{sl}}_q^n(2)$, respectively.

Throughout the chapter, all vector spaces are assumed to be vector spaces over the complex number field C and the following notations are used:

- $C^* := \{x \in C | x \neq 0\}$.
- q is a complex number satisfying $q^2 \neq 0, 1$.
- $\alpha(q) := t$ if q is a primitive t -th root of unity.
-

$$T := \begin{cases} \alpha(q), & \text{if } \alpha(q) \text{ is odd,} \\ \frac{\alpha(q)}{2}, & \text{if } \alpha(q) \text{ is even.} \end{cases}$$

-

$$[m] := \frac{q^m - q^{-m}}{q - q^{-1}}, \quad \langle m \rangle := \frac{q^m + q^{-m}}{2}, \quad \text{where } m \in \mathbb{Z}.$$

- For any $\mathbb{Z}$ -graded vector space $V = \bigoplus_{n \in \mathbb{Z}} V_n$, $J \in \text{Hom}(V, V)$ is defined by $J(v_n) := q^n v_n$ for $n \in \mathbb{Z}$ and $v_n \in V_n$.
- $\sigma \in \text{Hom}(V, V)$ is defined by $\sigma := \frac{J + J^{-1}}{2}$.

§1.1. J -derivations and the quantum Witt algebra

Definition 1.1 Let A be a $\mathbb{Z}$ -graded algebra. A linear map $D : A \rightarrow A$ is called a J -derivation if $D(ab) = D(a)J(b) + J^{-1}(a)D(b)$ for all $a, b \in A$.

Define a family of linear operators D_n on the Laurent polynomial ring $\mathbb{C}[t, t^{-1}]$ by

$$D_n := -t^{n+1} \left(\frac{d}{dt} \right)_q, \quad \left(\frac{d}{dt} \right)_q : t^m \mapsto [m]t^{m-1},$$

for all $m, n \in \mathbb{Z}$. Set $\mathcal{D} := \bigoplus_{n \in \mathbb{Z}} \mathbb{C}D_n$, then $\mathcal{D}$ is a $\mathbb{Z}$ -graded vector space with the $\mathbb{Z}$ -grading $\mathcal{D}_n := \mathbb{C}D_n$.

PROPOSITION 1.1. *With the notations above, we have*

- (1) $\mathcal{D}$ is the vector space formed by all J -derivations of $\mathbb{C}[t, t^{-1}]$.
- (2) The bracket $[,]$ on $\mathcal{D}$, defined by

$$[D_m, D_n] := JD_m J^{-1} D_n J - JD_n J^{-1} D_m J = [m - n]D_{m+n},$$

satisfies the following the **quantum Jacobi identity**:

$$(1.1) \quad [[x, y], \sigma(z)] + [[y, z], \sigma(x)] + [[z, x], \sigma(y)] = 0$$

where $x, y, z \in \mathcal{D}$.

PROOF: (1). For $m, n, k \in \mathbb{Z}$, we have

$$\begin{aligned} D_n(t^m \cdot t^k) &= -t^{n+1} \left(\frac{d}{dt} \right)_q (t^{m+k}) = -t^{n+1} [m+k] t^{m+k-1} \\ &= -([m]q^k + [k]q^{-m}) t^{n+m+k} \\ &= (-t^{n+1} [m] t^{m-1}) (q^k t^k) + (q^{-m} t^m) (-t^{n+1} [k] t^{k-1}) \\ &= D_n(t^m) J(t^k) + J^{-1}(t^m) D_n(t^k), \end{aligned}$$

so D_n is a J -derivation of $\mathbb{C}[t, t^{-1}]$. Conversely, if D is a J -derivation of $\mathbb{C}[t, t^{-1}]$, then $D(t) \in \mathbb{C}[t, t^{-1}]$. Using induction on m , we have that

$$D(t^m) = [m]t^{m-1} D(t) = D(t) \left(\frac{d}{dt} \right)_q (t^m) \quad \text{for } m \in \mathbb{Z}.$$

Hence, $D = D(t) \left(\frac{d}{dt} \right)_q \in \bigoplus_{n \in \mathbb{Z}} \mathbb{C}D_n = \mathcal{D}$.

(2). For $k \in \mathbb{Z}$ and $t^k \in \mathbb{C}[t, t^{-1}]$, we have

$$\begin{aligned} JD_m J^{-1} D_n J(t^k) &= -q^{m+k} [n+k] D_{m+n}(t^k), \\ JD_n J^{-1} D_m J(t^k) &= -q^{n+k} [m+k] D_{m+n}(t^k), \end{aligned}$$

so

$$\begin{aligned}
& (JD_m J^{-1} D_n J - JD_n J^{-1} D_m J)(t^k) \\
&= (q^{n+k}[m+k] - q^{m+k}[n+k])D_{m+n}(t^k) \\
&= [m-n]D_{m+n}(t^k),
\end{aligned}$$

or $JD_m J^{-1} D_n J - JD_n J^{-1} D_m J = [m-n]D_{m+n}$. In order to prove that the bracket $[,]$ on $\mathcal{D}$ satisfies the quantum Jacobi identity (1.1), it suffices to prove the following:

$$\begin{aligned}
(1.2) \quad & \langle k \rangle [[D_m, D_n], D_k] + \langle m \rangle [[D_n, D_k], D_m] \\
& + \langle n \rangle [[D_k, D_m], D_n] = 0,
\end{aligned}$$

where $m, n, k \in \mathbb{Z}$. Using $[D_m, D_n] = [m-n]D_{m+n}$, we get

$$\begin{aligned}
& \langle k \rangle [[D_m, D_n], D_k] \\
&= \langle k \rangle [m-n][D_{m+n}, D_k] = \langle k \rangle [m-n][m+n-k]D_{m+n+k} \\
&= (q^k + q^{-k})(q^{2m-k} + q^{k-2m} - q^{2n-k} - q^{k-2n}) \frac{D_{m+n+k}}{2(q-q^{-1})^2} \\
&= ((q^{2m} + q^{-2m}) + (q^{2k-2m} + q^{2m-2k}) \\
&\quad - (q^{2n} + q^{-2n}) - (q^{2n-2k} + q^{2k-2n})) \frac{D_{m+n+k}}{2(q-q^{-1})^2},
\end{aligned}$$

or

$$\begin{aligned}
(1.3) \quad & \langle k \rangle [[D_m, D_n], D_k] \\
&= (\langle 2m \rangle + \langle 2m-2k \rangle - \langle 2n \rangle - \langle 2n-2k \rangle) \frac{D_{m+n+k}}{(q-q^{-1})^2}.
\end{aligned}$$

Replacing m, n and k by n, k and m in (1.3) respectively, we get

$$\begin{aligned}
(1.4) \quad & \langle m \rangle [[D_n, D_k], D_m] \\
&= (\langle 2n \rangle + \langle 2n-2m \rangle - \langle 2k \rangle - \langle 2k-2m \rangle) \frac{D_{m+n+k}}{(q-q^{-1})^2}.
\end{aligned}$$

Replacing m, n and k by n, k and m in (1.4) respectively, we get

$$\begin{aligned}
(1.5) \quad & \langle n \rangle [[D_k, D_m], D_n] \\
&= (\langle 2k \rangle + \langle 2k-2n \rangle - \langle 2m \rangle - \langle 2m-2n \rangle) \frac{D_{m+n+k}}{(q-q^{-1})^2}.
\end{aligned}$$

Now (1.2) follows from (1.3) + (1.4) + (1.5). ■

Definition 1.2 A $\mathbb{Z}$ -graded vector space $L = \bigoplus_{n \in \mathbb{Z}} L_n$ is called a **quantum Lie algebra** if there is a skew-symmetric bilinear map $[\cdot, \cdot] : L \times L \rightarrow L$ which satisfies the quantum Jacobi identity (1.1) for all $x, y, z \in L$ and preserves $\mathbb{Z}$ -grading, i.e. $[L_m, L_n] \subseteq L_{m+n}$ for $m, n \in \mathbb{Z}$.

Let $L = \bigoplus_{n \in \mathbb{Z}} L_n$ be a quantum Lie algebra. The set $L(\mathbb{Z}) := \{n \in \mathbb{Z} | L_n \neq 0\}$ is called the **degree set** of L . A **graded subalgebra** (resp. a **graded ideal**) of L is a subalgebra (resp. an ideal) of L which is, in addition, a $\mathbb{Z}$ -graded subspace of the $\mathbb{Z}$ -graded vector space L .

Let $L = \bigoplus_{n \in \mathbb{Z}} L_n$ and $L' = \bigoplus_{n \in \mathbb{Z}} L'_n$ be quantum Lie algebras. A linear map $f : L \rightarrow L'$ is called a **homomorphism of type r** if $f([x, y]) = [f(x), f(y)]$ for $x, y \in L$ and there exists a rational number r such that $rL(\mathbb{Z}) \subseteq \mathbb{Z}$, $f(L_n) \subseteq L'_{rn}$ for $n \in L(\mathbb{Z})$. A homomorphism f of type r is called an **isomorphism of type r** if f is a bijection. A homomorphism of type 1 is also called **$\mathbb{Z}$ -grading preserving homomorphism**.

Example 1. By Proposition 1.1(2), the $\mathbb{Z}$ -graded vector space $W_q := \bigoplus_{n \in \mathbb{Z}} \mathbb{C}d_n$ with the $\mathbb{Z}$ -grading $(W_q)_n := \mathbb{C}d_n$ becomes a quantum Lie algebra under the following bracket:

$$[d_m, d_n] := [m - n]d_{m+n}, \quad \text{where } m, n \in \mathbb{Z},$$

which is called the **quantum Witt algebra**.

For a fixed positive integer n , we use $sl_q^n(2)$ to denote the three dimensional graded subalgebra $\mathbb{C}d_{-n} \oplus \mathbb{C}d_0 \oplus \mathbb{C}d_n$ of W_q . Note that $sl_q^n(2)$ has the $\mathbb{Z}$ -grading:

$$(sl_q^n(2))_k := \mathbb{C}d_k, \quad \text{where } k = 0, \pm n.$$

If $q^{4n} \neq 1$ and $\lambda \in \mathbb{C}^*$, then the linear map f defined by

$$f(d_n) := \frac{[2n][n]}{[2]} \lambda^{-1} d'_1, \quad f(d_{-n}) := \lambda d'_{-1}, \quad f(d_0) := [n]d'_0$$

is an isomorphism of type $\frac{1}{n}$ from $sl_q^n(2)$ to $sl_q^1(2) := \mathbb{C}d'_{-1} \oplus \mathbb{C}d'_0 \oplus \mathbb{C}d'_1$, where $[d'_i, d'_j] := [i - j]d'_{i+j}$ for $i, j \in \{0, \pm 1\}$.

Example 2. Let q be not a root of unity. We define the **quantum Virasoro algebra** V_q by

$$V_q := \bigoplus_{n \in \mathbb{Z}} \mathbb{C}d_n \oplus \mathbb{C}c,$$

$$(V_q)_n := \begin{cases} \mathbb{C}d_0 \oplus \mathbb{C}c, & \text{if } n = 0, \\ \mathbb{C}d_n, & \text{if } n \neq 0, \end{cases}$$

$$[d_m, d_n] := [m - n]d_{m+n} + \frac{[m-1][m][m+1]}{[2][3] \langle m \rangle} \delta_{m+n,0} c,$$

where $m, n \in \mathbb{Z}$. One can check directly that V_q is a quantum Lie algebra.

Example 3. If A is an associative algebra and $L = \bigoplus_{n \in \mathbb{Z}} L_n$ a quantum Lie algebra, then $A \otimes L$ becomes a quantum Lie algebra under the bracket defined by

$$[a \otimes x, b \otimes y] := ab \otimes [x, y], \quad \text{where } a, b \in A, x, y \in L,$$

and the $\mathbb{Z}$ -grading: $(A \otimes L)_n := A \otimes L_n$ for $n \in \mathbb{Z}$. In particular, $\mathbb{C}[t, t^{-1}] \otimes s\ell_q^n(2)$ is a quantum Lie algebra.

Example 4. Let n be a fixed positive integer. If $q^{4n} \neq 1$, then we define the simplest **quantum Kac-Moody algebra** $s\widehat{\ell}_q^n(2)$ as follows:

$$s\widehat{\ell}_q^n(2) := \mathbb{C}[t, t^{-1}] \otimes s\ell_q^n(2) \oplus \mathbb{C}c,$$

$$\left(s\widehat{\ell}_q^n(2)\right)_k := \begin{cases} 0, & \text{if } k \neq 0, \pm n, \\ \mathbb{C}[t, t^{-1}] \otimes \mathbb{C}d_k, & \text{if } k = \pm n, \\ \mathbb{C}[t, t^{-1}] \otimes \mathbb{C}d_0 \oplus \mathbb{C}c, & \text{if } k = 0, \end{cases}$$

$$[u \otimes d_0, v \otimes d_0] := -\text{Res} \left(u \frac{dv}{du} \right) c,$$

$$[u \otimes d_0, v \otimes d_{\pm n}] := \mp [n]uv \otimes d_{\pm n},$$

$$[u \otimes d_n, v \otimes d_{-n}] := [2n]uv \otimes d_0 + \frac{[2n]}{[n] \langle n \rangle} \text{Res} \left(u \frac{dv}{dt} \right) c,$$

$$[u \otimes d_n, v \otimes d_n] := [u \otimes d_{-n}, v \otimes d_{-n}] := [c, s\widehat{\ell}_q^n(2)] := 0,$$

where $u, v \in \mathbb{C}[t, t^{-1}]$, Res and $\frac{d}{dt}$ are the usual residue and derivation, respectively. Using the following well-known fact:

$$\text{Res} \left(uv \frac{dw}{dt} \right) + \text{Res} \left(vw \frac{du}{dt} \right) + \text{Res} \left(wu \frac{dv}{dt} \right) = 0,$$

where $u, v, w \in \mathbb{C}[t, t^{-1}]$, it is easy to check that $s\widehat{\ell}_q^n(2)$ is a quantum Lie algebra.

A quantum Lie algebra L is called **graded simple** (resp. **simple**) if $[L, L] \neq 0$ and L does not have any graded ideals (resp. ideals) which are different from $\{0\}$ and L .

PROPOSITION 1.2. For quantum Witt algebra W_q , we have

- (1) W_q is graded simple $\iff q$ is not a root of unity or $o(q) \neq 4$.
- (2) W_q is simple $\iff q$ is not a root of unity.

PROOF: (1). $\Leftarrow$: Let $0 \neq N$ be a graded ideal of W_q . We have two cases to discuss.

• If q is not a root of unity, then

$$d_m \in N \quad \text{for some } m \in \mathbb{Z}.$$

Since $N \ni [d_m, d_{-m}] = [2m]d_0$, we have

$$(1.6) \quad d_0 \in N.$$

Therefore, $N \ni [d_k, d_0] = [k]d_k$, for $k \in \mathbb{Z}$. This implies that

$$(1.7) \quad d_k \in N \quad \text{for } k \in \mathbb{Z} \setminus \{0\}.$$

It follows from (1.6) and (1.7) that $N = W_q$.

• If q is a root of unity and $o(q) \neq 4$, then $[2] \neq 0$ and

$$d_{kT+i} \in N \quad \text{for some } k \in \mathbb{Z} \text{ and } 0 \leq i \leq T-1.$$

If $2i \notin T\mathbb{Z}$, then $[2i] \neq 0$ and

$$N \ni [d_{kT+i}, d_{-kT-i}] = [2kT+2i]d_0 = [2i]d_0,$$

which implies (1.6).

If $2i \in T\mathbb{Z}$, then $[2i+1] \neq 0$ and

$$N \ni [d_{kT+i}, d_{-kT-i}] = [2kT+2i+1]d_1 = [2i+1]d_1.$$

So

$$d_1 \in N \quad \text{and} \quad N \ni [d_1, d_{-1}] = [2]d_0,$$

which also implies (1.6).

Hence, we proved that (1.6) always holds.

In order to prove that $N = W_q$, it suffices to prove

$$(1.8) \quad d_{mT+r} \in N \quad \text{for } m \in \mathbb{Z} \text{ and } 0 \leq r \leq T-1$$

If $r \neq 0$, then, by (1.6)

$$N \ni [d_{mT+r}, d_0] = [mT+r]d_{mT+r} = q^{mT}[r]d_{mT+r}.$$

So

$$(1.9) \quad d_{mT+r} \in N \quad \text{for } m \in \mathbb{Z} \text{ and } 1 \leq r \leq T-1.$$

If $r = 0$, then $d_{mT+1} \in N$ by (1.9). Hence,

$$N \ni [d_{mT+1}, d_{-1}] = [mT + 2]d_{mT} = q^{mT}[2]d_{mT},$$

i.e., $d_{mT} \in N$ for $m \in \mathbb{Z}$.

(1). $\Rightarrow$: If $o(q) = 4$, then $N := \bigoplus_{n \in \mathbb{Z}} \mathbb{C}d_{2n+1}$ is a graded ideal of W_q , so W_q is not graded simple.

(2). $\Leftarrow$: Let $0 \neq N$ be an ideal of W_q and let

$$(1.10) \quad 0 \neq x = a_1 d_{m_1} + a_2 d_{m_2} + \cdots + a_\ell d_{m_\ell} \in N,$$

where $a_i \in \mathbb{C}^*$, $m_i \in \mathbb{Z}$ and $m_i \neq m_j$ if $1 \leq i \neq j \leq \ell$.

We choose $x \in N \setminus \{0\}$ such that ℓ in (1.10) is minimal. Then $\ell = 1$. In fact, if $\ell \geq 2$, we have

$$\begin{aligned} N \ni [d_{m_1}, x] &= a_2 [m_1 - m_2] d_{m_2} + \cdots + a_\ell [m_1 - m_\ell] d_{m_\ell}, \\ a_i [m_1 - m_i] &\neq 0 \quad \text{for } 2 \leq i \leq \ell, \end{aligned}$$

which contradicts to the choice of ℓ .

Therefore, we have proved that there exists some $d_m \in N$. As above, this implies that $N = W_q$.

(2). $\Rightarrow$: If q is a root of unity, then $N(a_i, k_i, 0 \leq i \leq \ell)$ defined by

$$N(a_i, k_i, 0 \leq i \leq \ell) := \bigoplus_{\substack{k \in \mathbb{Z} \\ 0 \leq s \leq T-1}} \mathbb{C} \left(\sum_{i=0}^{\ell} a_i d_{(k_i+k)T+s} \right)$$

is an ideal which is different from $\{0\}$ and W_q , where ℓ is a positive integer, $a_0, a_1, \dots, a_\ell \in \mathbb{C}^*$, $k_0, k_1, \dots, k_\ell \in \mathbb{Z}$, $k_i - k_j \in 2\mathbb{Z}$ and $k_i \neq k_j$, for $0 \leq i < j \leq \ell$. Therefore, W_q is not simple. ■

§1.2. Quantum central extensions

In this section, we assume that all homomorphisms between quantum Lie algebras are $\mathbb{Z}$ -grading preserving homomorphisms.

An exact sequence of quantum Lie algebras

$$(2.1) \quad 0 \longrightarrow \mathcal{C} \xrightarrow{i} E \xrightarrow{\pi} L \longrightarrow 0$$

is said to be a **quantum extension** of L by C , which is denoted by $[i, \pi]$. This quantum extension is said to be a **quantum central extension** if $[i(C), E] = 0$. A **morphism** of the quantum extension (2.1) to another extension

$$(2.2) \quad 0 \longrightarrow C' \xrightarrow{i'} E' \xrightarrow{\pi'} L \longrightarrow 0$$

is a pair (ψ, φ) of quantum Lie algebra homomorphisms such that the diagram

$$\begin{array}{ccccc} C & \xrightarrow{i} & E & \xrightarrow{\pi} & L \\ \psi \downarrow & & \varphi \downarrow & & \parallel \\ C' & \xrightarrow{i'} & E' & \xrightarrow{\pi'} & L \end{array}$$

is commutative. A quantum central extension (2.1) of L is called **quantum universal central extension** if, given any quantum central extension (2.2) of L , there is a unique morphism from (2.1) to (2.2). Two quantum extensions E and E' of L by C are **equivalent** if there is an isomorphism $\varphi : E \simeq E'$ making the following diagram commutes:

$$\begin{array}{ccccccc} 0 & \longrightarrow & C & \longrightarrow & E & \longrightarrow & L \longrightarrow 0 \\ & & \parallel & & \varphi \downarrow & & \parallel \\ 0 & \longrightarrow & C & \longrightarrow & E' & \longrightarrow & L \longrightarrow 0 \end{array}$$

Let V be a $\mathbb{Z}$ -graded vector space and L a quantum Lie algebra. A skew-symmetric bilinear maps $\alpha : L \times L \rightarrow V$ is said to be a **quantum 2-cocycle** of L with coefficients in V if α satisfies the following conditions:

$$(2.3) \quad \alpha(L_n, L_m) \subseteq V_{n+m},$$

$$(2.4) \quad \alpha([x, y], \sigma(z)) + \alpha([y, z], \sigma(x)) + \alpha([z, x], \sigma(y)) = 0$$

for all $n, m \in \mathbb{Z}$ and $x, y, z \in L$.

Let $Z_q^2(L, V)$ be the space of all quantum 2-cocycles of L with coefficients in V and $B_q^2(L, V)$ the space of all skew-symmetric bilinear map $\alpha : L \times L \rightarrow V$ satisfying (2.3) and

$$(2.5) \quad \alpha(x, y) = f([x, y])$$

for $x, y \in L$ and some $\mathbb{Z}$ -grading preserving linear map $f : L \rightarrow V$. It is clear that $B_q^2(L, V) \subseteq Z_q^2(L, V)$. The quotient $H_q^2(L, V) := \frac{Z_q^2(L, V)}{B_q^2(L, V)}$ is called the **quantum second cohomology group** of L with coefficients in V .

Let (2.1) be a quantum central extension of L by C , where $[C, C] = 0$. Then the **equivalent class** $[[i, \pi]]$ containing the quantum central extension (2.1) is defined to be the set of all quantum central extensions of L by C which are equivalent to (2.1). Let $CE_q(L, C)$ denote the set of all equivalent classes of quantum central extensions of L by C .

Thus, if L and C are quantum Lie algebras with $[C, C] = 0$, then

$$H_q^2(L, C) = \{ [\alpha] := \alpha + B_q^2(L, C) | \alpha \in Z_q^2(L, C) \},$$

$$CE_q(L, C) = \left\{ [[i, \pi]] \mid \begin{array}{l} [i, \pi] \text{ is a quantum central exten-} \\ \text{sion of } L \text{ by } C \end{array} \right\}.$$

Now we construct two maps F and G between $H_q^2(L, C)$ and $CE_q(L, C)$ as follows:

• $F : H_q^2(L, C) \rightarrow CE_q(L, C)$ is defined by $F([\alpha]) = [[i_\alpha, \pi_\alpha]]$ where

$$(2.6) \quad 0 \longrightarrow C \xrightarrow{i_\alpha} E_\alpha \xrightarrow{\pi_\alpha} L \longrightarrow 0$$

is defined by

$$E_\alpha := \oplus_{n \in \mathbb{Z}} (E_\alpha)_n, \quad \text{where } (E_\alpha)_n := L_n \oplus C_n,$$

$$[(x, u), (y, v)] := ([x, y], \alpha(x, y)),$$

$$i_\alpha : u \mapsto (0, u), \quad \pi_\alpha : (x, u) \mapsto x$$

for $x, y \in L$ and $u, v \in C$.

• $G : CE_q(L, C) \rightarrow H_q^2(L, C)$ is defined by

$$G([[i, \pi]]) = [\alpha], \quad \alpha_0 = i\alpha,$$

$$\alpha_0(x, y) := [\beta(x), \beta(y)] - \beta([x, y]) \quad \text{for } x, y \in L,$$

where $\beta : L \rightarrow E$ is a $\mathbb{Z}$ -grading preserving linear map satisfying $\pi\beta = id|_L$.

Using the notations above, we have

PROPOSITION 2.1. *The map F is a bijection with inverse G .*

PROOF: What we need to check is the following:

- (i) F is well-defined;
- (ii) G is well-defined;
- (iii) $FG = 1$ and $GF = 1$.

(i). For $x, y, z \in L$ and $u, v, w \in \mathcal{C}$, we have

$$\begin{aligned}
& [[(x, u), (y, v)], \sigma((z, w))] + [[(y, v), (z, w)], \sigma((x, u))] \\
& \quad + [[(z, w), (x, u)], \sigma((y, v))] \\
& = [[(x, y), \alpha(x, y)], (\sigma(z), \sigma(w))] + [[(y, z), \alpha(y, z)], (\sigma(x), \sigma(u))] \\
& \quad + [[(z, x), \alpha(z, x)], (\sigma(y), \sigma(v))] \\
& = ([[x, y], \sigma(z)], \alpha([x, y], \sigma(z))) + ([[y, z], \sigma(x)], \alpha([y, z], \sigma(x))) \\
& \quad + ([[z, x], \sigma(y)], \alpha([z, x], \sigma(y))) \\
& = 0
\end{aligned}$$

Also, it is easy to check that

$$\begin{aligned}
& [(E_\alpha)_m, (E_\alpha)_n] \subseteq (E_\alpha)_{m+n} \quad \text{for } m, n \in \mathbb{Z}, \\
& [(x, u), (y, v)] = -[(y, v), (x, u)] \quad \text{for } x, y \in L \text{ and } u, v \in \mathcal{C}.
\end{aligned}$$

Therefore, E_α is a quantum Lie algebra. It is obvious that i_α is injective and π_α is surjective. So we have proved that (2.6) is a quantum central extension of L by $\mathcal{C}$.

Next, we prove that (2.6) is independent of the choice of α .

Suppose that α and $\alpha' \in Z_q^2(L, \mathcal{C})$ and $\alpha' - \alpha \in B_q^2(L, \mathcal{C})$, i.e., there exists a $\mathbb{Z}$ -grading preserving linear map $f : L \rightarrow \mathcal{C}$ such that

$$(\alpha' - \alpha)(x, y) = f([x, y]) \quad \text{for } x, y \in L.$$

We define $\varphi : E_\alpha \rightarrow E_{\alpha'}$ by

$$\varphi : (x, u) \mapsto (x, u + f(x)), \quad \text{where } x \in L \text{ and } u \in \mathcal{C}.$$

On one hand, if we denote the bracket on E_α and $E_{\alpha'}$ by $[\cdot, \cdot]_\alpha$ and $[\cdot, \cdot]_{\alpha'}$, respectively, then, for $x, y \in L$ and $u, v \in \mathcal{C}$,

$$\begin{aligned}
\varphi([(x, u), (y, v)]_\alpha) &= \varphi([x, y], \alpha(x, y)) \\
&= ([x, y], \alpha(x, y) + f([x, y])) \\
&= ([x, y], \alpha'(x, y)) \\
&= [(x, u), (y, v)]_{\alpha'} \\
&= [(x, u + f(x)), (y, v + f(y))]_{\alpha'} \\
&= [\varphi((x, u)), \varphi((y, v))]_{\alpha'}.
\end{aligned}$$

It follows that φ is an isomorphism from the quantum Lie algebra E_α to the quantum Lie algebra $E_{\alpha'}$.

On the other hand, φ makes the following diagram commutative:

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{C} & \xrightarrow{i_\alpha} & E_\alpha & \xrightarrow{\pi_\alpha} & L \longrightarrow 0 \\ & & \parallel & & \varphi \downarrow & & \parallel \\ 0 & \longrightarrow & \mathcal{C} & \xrightarrow{i_{\alpha'}} & E_{\alpha'} & \xrightarrow{\pi_{\alpha'}} & L \longrightarrow 0 \end{array}$$

Therefore, $[i_\alpha, \pi_\alpha]$ and $[i_{\alpha'}, \pi_{\alpha'}]$ are equivalent.

This proves (i).

(ii). Fix a $\mathbb{Z}$ -grading preserving linear map $\beta : L \rightarrow E$ which satisfies $\pi\beta = id|_L$. Define $\alpha_0 : L \times L \rightarrow E$ by

$$(2.7) \quad \alpha_0(x, y) := [\beta(x), \beta(y)] - \beta([x, y]), \quad \text{for } x, y \in L.$$

Then $\alpha_0(x, y) \in \text{Ker}(\pi) = \text{Im}(i)$. Using the injectivity of i , we can construct $\alpha : L \times L \rightarrow \mathcal{C}$ such that $\alpha_0 = i\alpha$. For $x, y, z \in L$,

$$\begin{aligned} & \alpha_0([x, y], \sigma(z)) + \alpha_0([y, z], \sigma(x)) + \alpha_0([z, x], \sigma(y)) \\ &= ([[\beta(x), \beta(y)], \sigma(\beta(z))]) + ([[\beta(y), \beta(z)], \sigma(\beta(x))]) \\ & \quad + ([[\beta(z), \beta(x)], \sigma(\beta(y))]) \\ & \quad - \beta([[\beta(x), \sigma(z)] + [[\beta(y), \sigma(x)] + [[\beta(z), \sigma(y)]] = 0, \end{aligned}$$

by (2.7) and the fact: $\sigma\beta = \beta\sigma$. This implies that α is a quantum 2-cocycle.

Now we prove that the equivalent class $[\alpha]$ constructed from β is independent of the choice of β . Suppose $\beta' : L \rightarrow E$ is another $\mathbb{Z}$ -grading preserving linear map satisfying $\pi\beta' = id|_L$. Using the same procedure as above, we can construct α'_0 and a quantum 2-cocycle α' such that

$$\alpha'_0 = i\alpha', \quad \alpha'_0(x, y) = [\beta'(x), \beta'(y)] - \beta'([x, y]),$$

where $x, y \in L$. That $\text{Im}(\beta' - \beta) \in \text{Ker}(\pi) = \text{Im}(i)$ produces a $\mathbb{Z}$ -grading preserving linear map $f : L \rightarrow \mathcal{C}$ satisfying $\beta' - \beta = if$. It follows that

$$\begin{aligned} i(\alpha - \alpha')(x, y) &= (\alpha_0 - \alpha'_0)(x, y) \\ &= [(\beta - \beta')(x), \beta(y)] + [\beta'(x), (\beta - \beta')(y)] + if([x, y]) = if([x, y]). \end{aligned}$$

Thus $(\alpha - \alpha')((x, y)) = f([x, y])$, i.e. $\alpha - \alpha' \in B_q^2(L, \mathcal{C})$, or $[\alpha] = [\alpha']$.

(iii). Let $[[i, \pi]] \in CE_q(L, \mathcal{C})$, then $FG([[i, \pi]]) = [[i_\alpha, \pi_\alpha]]$, where $[i, \pi]$ and $[i_\alpha, \pi_\alpha]$ denote the quantum central extensions (2.1) and (2.6), respectively.

In order to prove that $[i, \pi]$ and $[i_\alpha, \pi_\alpha]$ are equivalent, we define $\varphi : E_\alpha \rightarrow E$ by

$$\varphi((x, u)) := \beta(x) + i(u), \quad \text{where } x \in L \text{ and } u \in \mathcal{C}.$$

Then, for $x, y \in L$ and $u, v \in \mathcal{C}$, we have

$$\begin{aligned}
\varphi([(x, u), (y, v)]) &= \varphi([x, y], \alpha(x, y)) \\
&= \beta([x, y]) + i\alpha(x, y) \\
&= \beta([x, y]) + \alpha_0(x, y) = [\beta(x), \beta(y)] \\
&= [\beta(x) + i(u), \beta(y) + i(v)] \\
&= [\varphi((x, u)), \varphi((y, v))].
\end{aligned}$$

This proves that φ is a $\mathbb{Z}$ -grading preserving homomorphism.

Furthermore, the following diagram is commutative:

$$\begin{array}{ccccccc}
0 & \longrightarrow & \mathcal{C} & \xrightarrow{i_\alpha} & E_\alpha & \xrightarrow{\pi_\alpha} & L \longrightarrow 0 \\
& & \parallel & & \downarrow \varphi & & \parallel \\
0 & \longrightarrow & \mathcal{C} & \xrightarrow{i} & E & \xrightarrow{\pi} & L \longrightarrow 0
\end{array}$$

In fact, for $x \in L$ and $u \in \mathcal{C}$, we have

$$\varphi i_\alpha(u) = \varphi((0, u)) = \beta(0) + i(u) = i(u)$$

and

$$\pi \varphi((x, u)) = \pi(\beta(x) + i(u)) = x = \pi_\alpha((x, u)).$$

It follows that φ is a $\mathbb{Z}$ -grading preserving isomorphism. This means that $[i, \pi]$ and $[i_\alpha, \pi_\alpha]$ are equivalent, i.e., $[[i, \pi]] = [[i_\alpha, \pi_\alpha]]$.

Thus, we have proved that $FG = 1$.

Similarly, one can check $GF = 1$. ■

PROPOSITION 2.2. *Let L be a quantum Lie algebra. If $[L, L] = L$, then L has a quantum universal central extension.*

PROOF: Let $L = \bigoplus_{n \in \mathbb{Z}} L_n$ be a quantum Lie algebra. We define the q -analogue of the usual second homology group of L with trivial coefficients by $H_2^q(L, \mathbb{C}) := \frac{\Lambda^2(L)}{I}$, where I is the $\mathbb{Z}$ -graded subspace of $\Lambda^2(L)$ spanned by all elements of the form

$$[x, y] \wedge \sigma(z) + [y, z] \wedge \sigma(x) + [z, x] \wedge \sigma(y),$$

where $x, y, z \in L$.

Let $\alpha : L \times L \rightarrow H_2^q(L, \mathbb{C})$ be the map defined by

$$\alpha : (x, y) \mapsto x \wedge y + I \quad \text{for } x, y \in L.$$

Then $\alpha \in Z_q^2(L, H_2^q(L, \mathbb{C}))$. Let $F([\alpha])$ be the quantum central extension:

$$(2.8) \quad 0 \longrightarrow H_2^q(L, \mathbb{C}) \xrightarrow{i_\alpha} E_\alpha \xrightarrow{\pi_\alpha} L \longrightarrow 0$$

as constructed prior to Proposition 2.1.

Suppose that

$$(2.9) \quad 0 \longrightarrow \mathcal{C}' \xrightarrow{i'} E' \xrightarrow{\pi'} L \longrightarrow 0$$

is any quantum central extension of L . We denote the bracket on E' by $[\cdot, \cdot]'$. Let us write E' as a direct sum of subspaces $\oplus_{n \in \mathbb{Z}} (L'_n \oplus i'(\mathcal{C}'_n))$, where $L' := \oplus_{n \in \mathbb{Z}} L'_n$ is a preimage of L under π' . Identifying L with L' and $\mathcal{C}'$ with $i'(\mathcal{C}')$, we have $E' = \oplus_{n \in \mathbb{Z}} (E')_n$, where $(E')_n = L_n \oplus (\mathcal{C}')_n$.

Then we have a quantum 2-cocycle $\beta : L \times L \rightarrow \mathcal{C}'$ with

$$[x, y]' = [x, y] + \beta(x, y) \quad \text{for } x, y \in L.$$

It follows that there is an induced map $\beta : \wedge^2(L) \rightarrow \mathcal{C}'$ vanishing on the subspace $I \subseteq \wedge^2(L)$ defined above. This allows us to define a linear map $\psi : H_2^q(L, \mathbb{C}) \rightarrow \mathcal{C}'$ by

$$\psi(\alpha(x, y)) = \beta(x, y), \quad \text{where } x, y \in L.$$

Next, we define the linear map $\varphi : E_\alpha \rightarrow E'$ by

$$\varphi((x, u)) = x + \psi(u), \quad \text{where } x \in L \text{ and } u \in H_2^q(L, \mathbb{C}).$$

Then the following diagram is commutative:

$$\begin{array}{ccccc} H_2^q(L, \mathbb{C}) & \xrightarrow{i_\alpha} & E_\alpha & \xrightarrow{\pi_\alpha} & L \\ \psi \downarrow & & \varphi \downarrow & & \parallel \\ \mathcal{C}' & \xrightarrow{i'} & E' & \xrightarrow{\pi'} & L \end{array}$$

In fact, for $x \in L$ and $u \in H_2^q(L, \mathbb{C})$, we have

$$\pi' \varphi((x, u)) = \pi'(x + \psi(u)) = x = \pi_\alpha((x, u))$$

and

$$\varphi i_\alpha(\alpha(x, y)) = \varphi((0, \alpha(x, y))) = \psi \alpha(x, y) = i' \psi(\alpha(x, y)).$$

It follows that $\pi' \varphi = id \cdot \pi_\alpha$ and $\varphi i_\alpha = i' \psi$.

If $x, y \in L$ and $u, v \in H_2^q(L, \mathbb{C})$, then

$$\begin{aligned}\varphi([(x, u), (y, v)]) &= \varphi([(x, y], \alpha(x, y))) \\ &= [x, y] + \psi\alpha(x, y) \\ &= [x, y] + \beta(x, y) = [x, y]' \\ &= [x + \psi(u), y + \psi(v)]' = [\varphi((x, u)), \varphi((y, v))].\end{aligned}$$

So φ is a $\mathbb{Z}$ -grading preserving homomorphism.

This proves that (ψ, φ) is a morphism from (2.8) to (2.9).

Let $E := [E_\alpha, E_\alpha]$. Since $[L, L] = L$, $\pi_\alpha(E) = L$ and $E_\alpha = E + \text{Im}(i_\alpha)$. Thus

$$E = [E_\alpha, E_\alpha] = [E + \text{Im}(i_\alpha), E + \text{Im}(i_\alpha)] = [E, E].$$

Let $\mathcal{C} := E \cap \text{Im}(i_\alpha)$. We will prove that

$$0 \longrightarrow \mathcal{C} \xrightarrow{i} E \xrightarrow{\pi} L \longrightarrow 0$$

is the quantum universal central extension of L , where i is the imbedding map and π is the restriction of π_α to E .

Indeed, given any other quantum central extension (2.9) of L , we have the following commutative diagram by what we have proved:

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{C} & \xrightarrow{i} & E & \xrightarrow{\pi} & L \longrightarrow 0 \\ & & \psi_0 \downarrow & & \varphi_0 \downarrow & & \parallel \\ 0 & \longrightarrow & H_2^q(L, \mathbb{C}) & \xrightarrow{i_\alpha} & E_\alpha & \xrightarrow{\pi_\alpha} & L \longrightarrow 0 \\ & & \psi \downarrow & & \varphi \downarrow & & \parallel \\ 0 & \longrightarrow & \mathcal{C}' & \xrightarrow{i'} & E' & \xrightarrow{\pi'} & L \longrightarrow 0 \end{array}$$

where φ_0 is the imbedding maps and ψ_0 is defined by

$$\psi_0 : i_\alpha(u) \mapsto u \quad \text{for } i_\alpha(u) \in \mathcal{C} \text{ and } u \in H_2^q(L, \mathbb{C})$$

Hence, $(\psi\psi_0, \varphi\varphi_0)$ is a morphism from $[i, \pi]$ to $[i', \pi']$.

Suppose that (ψ', φ') is any other morphism from $[i, \pi]$ to $[i', \pi']$, i.e.,

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{C} & \xrightarrow{i} & E & \xrightarrow{\pi} & L \longrightarrow 0 \\ & & \psi' \downarrow & & \varphi' \downarrow & & \parallel \\ 0 & \longrightarrow & \mathcal{C}' & \xrightarrow{i'} & E' & \xrightarrow{\pi'} & L \longrightarrow 0 \end{array}$$

Then, for all $z \in E$, we have $\pi'(\varphi\varphi_0)(z) = \pi(z) = \pi'\varphi'(z)$ and

$$(\varphi\varphi_0)(z) - \varphi'(z) \in \text{Ker}(\pi') = i'(C').$$

Now let $x, y \in E$, then

$$\begin{aligned} (\varphi\varphi_0)([x, y]) &= [(\varphi\varphi_0)(x), (\varphi\varphi_0)(y)] \\ &= [((\varphi\varphi_0)(x) - \varphi'(x)) + \varphi'(x), ((\varphi\varphi_0)(y) - \varphi'(y)) + \varphi'(y)] \\ &= [\varphi'(x), \varphi'(y)] = \varphi'([x, y]). \end{aligned}$$

It follows from $E = [E, E]$ that

$$\varphi\varphi_0 = \varphi'$$

and

$$i'(\psi\psi_0) = (\varphi\varphi_0)i = \varphi i = i'\psi'$$

or

$$\psi\psi_0 = \psi'$$

because i' is injective.

Thus, we have proved that for any quantum central extension $[i', \pi']$ of L , there exists a unique morphism $(\psi\psi_0, \varphi\varphi_0)$ from $[i, \pi]$ to $[i', \pi']$. Therefore, $[i, \pi]$ is the quantum universal central extension of L . ■

§1.3. The quantum universal central extension of W_q and $C[t, t^{-1}] \otimes s\ell_q^n(2)$

In this section, we will study the quantum universal central extension of the quantum Witt algebra W_q and the quantum Lie algebra $C[t, t^{-1}] \otimes s\ell_q^n(2)$, respectively.

THEOREM 3.1. *If q is not a root of unity, then the quantum Virasoro algebra V_q is the quantum universal central extension of the quantum Witt algebra W_q .*

PROOF: Let

$$0 \longrightarrow C \longrightarrow L \longrightarrow W_q \longrightarrow 0$$

be an arbitrary quantum central extension of W_q with corresponding quantum 2-cocycle α . By (2.4), we have

$$\alpha([d_m, d_n], \sigma(d_k)) + \alpha([d_n, d_k], \sigma(d_m)) + \alpha([d_k, d_m], \sigma(d_n)) = 0,$$

or

$$(3.1) \quad \begin{aligned} & \langle k \rangle [m - n] \alpha(d_k, d_{m+n}) + \langle m \rangle [n - k] \alpha(d_m, d_{n+k}) \\ & + \langle n \rangle [k - m] \alpha(d_n, d_{k+m}) = 0 \end{aligned}$$

Letting $k = 0$ in (3.1), we get

$$(3.2) \quad [m - n] \alpha(d_0, d_{m+n}) + [m + n] \alpha(d_m, d_n) = 0$$

Setting $k = 1$ and $n = -m - 1$ in (3.1), we obtain

$$(3.3) \quad \begin{aligned} & \langle m + 1 \rangle [m - 1] \alpha(d_{m+1}, d_{-m-1}) \\ & = \langle 1 \rangle [-2m - 1] \alpha(d_1, d_{-1}) + \langle m \rangle [m + 2] \alpha(d_m, d_{-m}) \end{aligned}$$

Set

$$\ell_n := \begin{cases} d_n - \frac{1}{[n]} \alpha(d_0, d_n), & \text{for } n \neq 0, \\ d_0 + \frac{1}{[2]} \alpha(d_1, d_{-1}), & \text{for } n = 0. \end{cases}$$

On one hand, if $m + n \neq 0$, then

$$\begin{aligned} [\ell_m, \ell_n] &= [d_m, d_n] = [m - n] d_{m+n} + \alpha(d_m, d_n) \\ &= [m - n] \left(\ell_{m+n} + \frac{1}{[m + n]} \alpha(d_0, d_{m+n}) \right) + \alpha(d_m, d_n) \\ &= [m - n] \ell_{m+n} + \left(\frac{[m - n]}{[m + n]} \alpha(d_0, d_{m+n}) + \alpha(d_m, d_n) \right). \end{aligned}$$

By (3.2), we have

$$(3.4) \quad [\ell_m, \ell_n] = [m - n] \ell_{m+n} \quad \text{if } m + n \neq 0.$$

On the other hand,

$$\begin{aligned} [\ell_m, \ell_{-m}] &= [d_m, d_{-m}] = [2m] d_0 + \alpha(d_m, d_{-m}) \\ &= [2m] \left(\ell_0 - \frac{1}{[2]} \alpha(d_1, d_{-1}) \right) + \alpha(d_m, d_{-m}) \\ (3.5) \quad &= [2m] \ell_0 + \left(-\frac{[2m]}{[2]} \alpha(d_1, d_{-1}) + \alpha(d_m, d_{-m}) \right). \end{aligned}$$

Now we define $\beta : W_q \times W_q \rightarrow \mathcal{C}$ by

$$\beta(d_m, d_n) = \begin{cases} 0, & \text{if } m + n \neq 0; \\ -\frac{[2m]}{[2]} \alpha(d_1, d_{-1}) + \alpha(d_m, d_{-m}), & \text{if } m + n = 0. \end{cases}$$

It follows from (3.4) and (3.5) that

$$(3.6) \quad [\ell_m, \ell_n] = [m-n]\ell_{m+n} + \beta(d_m, d_n).$$

Note β is a quantum 2-cocycle with $\beta(d_1, d_{-1}) = 0$. Hence, by (3.3)

$$< m+1 > [m-1]\beta(d_{m+1}, d_{-m-1}) = < m > [m+2]\beta(d_m, d_{-m}).$$

This implies that

$$\beta(d_m, d_{-m}) = \frac{[m-1][m][m+1]}{[2][3] < m >} < 2 > \beta(d_2, d_{-2}), \quad m \geq 2.$$

Setting $c = < 2 > \beta(d_2, d_{-2})$, (3.6) becomes

$$[\ell_m, \ell_n] = [m-n]\ell_{m+n} + \frac{[m-1][m][m+1]}{[2][3] < m >} \delta_{m+n,0} c,$$

which proves that V_q is the quantum universal central extension of W_q . ■

THEOREM 3.2. *If $q^{4n} \neq 1$, then $s\widehat{\ell}_q^n(2)$ is the quantum universal central extension of $\mathbb{C}[t, t^{-1}] \otimes s\ell_q^n(2)$.*

PROOF: Set $ux := u \otimes x$ for $u \in \mathbb{C}[t, t^{-1}]$ and $x \in s\ell_q^n(2)$. Let

$$0 \longrightarrow \mathcal{C} \longrightarrow L \xrightarrow{\pi} \mathbb{C}[t, t^{-1}] \otimes s\ell_q^n(2) \longrightarrow 0$$

be an arbitrary quantum central extension of $\mathbb{C}[t, t^{-1}] \otimes s\ell_q^n(2)$ and $a' \in L$ such that $\pi(a') = a$, then $[a', b']$ is independent of the choice of the preimages of a and b , where $a, b \in \mathbb{C}[t, t^{-1}] \otimes s\ell_q^n(2)$.

For $u \in \mathbb{C}[t, t^{-1}]$, we define $ud'_{\pm n}$ and ud'_0 in L by

$$(3.7) \quad ud'_{\pm n} := \mp \frac{1}{[n]} [d'_0, (ud_{\pm n})'], \quad ud'_0 := \frac{1}{[2n]} [d'_n, (ud_{-n})']$$

Set

$$(3.8) \quad \{u, v\} := [(ud_0)', (vd_0)'] \in \mathcal{C}, \quad u, v \in \mathbb{C}[t, t^{-1}]$$

Then we have

$$\pi(ud'_0) = \frac{1}{[2n]} [\pi(d'_n), \pi((ud_{-n})')] = \frac{1}{[2n]} [d_n, ud_{-n}] = ud_0,$$

and

$$\pi(ud'_{\pm n}) = \mp \frac{1}{[n]} [\pi(d'_0), \pi((ud_{\pm n})')] = \mp \frac{1}{[n]} [d_0, ud_{\pm n}] = ud_{\pm n}.$$

It follows that

$$(3.9) \quad [ud'_0, vd'_0] = \{u, v\},$$

$$(3.10) \quad ud'_{\pm n} = \mp \frac{1}{[n]} [d'_0, ud'_{\pm n}].$$

Since

$$\begin{aligned} \pi([vd'_0, ud'_{\pm n}]) &= [\pi(vd'_0), \pi(ud'_{\pm n})] \\ &= [vd_0, ud_{\pm n}] = \mp [n] uvd_{\pm n}, \end{aligned}$$

so we get

$$(3.11) \quad [vd'_0, ud'_{\pm n}] - (\mp [n] uvd'_{\pm n}) \in \mathcal{C}.$$

From

$$[vd'_0, [d'_0, ud'_{\pm n}]] + [d'_0, [ud'_{\pm n}, vd'_0]] + \langle n \rangle [ud'_{\pm n}, [vd'_0, d'_0]] = 0,$$

we get, by (3.8) and (3.9)

$$(3.12) \quad [vd'_0, [d'_0, ud'_{\pm n}]] = [d'_0, [vd'_0, ud'_{\pm n}]].$$

Using (3.10), (3.11) and (3.12),

$$\begin{aligned} [vd'_0, ud'_{\pm n}] &= [vd'_0, \mp \frac{1}{[n]} [d'_0, ud'_{\pm n}]] \\ &= \mp \frac{1}{[n]} [vd'_0, [d'_0, ud'_{\pm n}]] = \mp \frac{1}{[n]} [d'_0, [vd'_0, ud'_{\pm n}]] \\ &= \mp \frac{1}{[n]} [d'_0, \mp [n] uvd'_{\pm n}] = [d'_0, uvd'_{\pm n}] = \mp [n] uvd'_{\pm n}. \end{aligned}$$

This proves that

$$(3.13) \quad [vd'_0, ud'_{\pm n}] = \mp [n] uvd'_{\pm n}.$$

In particular, let $u = 1$ in (3.13), we have

$$(3.14) \quad [vd'_0, d'_{\pm n}] = \mp [n] vd'_{\pm n}.$$

From

$$\langle n \rangle [[ud'_0, d'_n], vd'_{-n}] + [[d'_n, vd'_{-n}], ud'_0] + \langle n \rangle [[vd'_{-n}, ud'_0], d'_n] = 0$$

and

$$(3.15) \quad [d'_n, vd'_{-n}] = [2n]vd'_0,$$

we get

$$(3.16) \quad [[ud'_0, d'_n], vd'_{-n}] = [[ud'_0, vd'_{-n}], d'_n] + \frac{[2n]}{\langle n \rangle} [ud'_0, vd'_0].$$

So, by (3.9), (3.13), (3.14) and (3.16)

$$\begin{aligned} [ud'_n, vd'_{-n}] &= [-\frac{1}{[n]} [ud'_0, d'_n], vd'_{-n}] \\ &= -\frac{1}{[n]} [[ud'_0, d'_n], vd'_{-n}] \\ &= -\frac{1}{[n]} [[ud'_0, vd'_{-n}], d'_n] - \frac{[2n]}{[n]\langle n \rangle} [ud'_0, vd'_0] \\ &= -\frac{1}{[n]} [[n]uvd'_{-n}, d'_n] - \frac{[2n]}{[n]\langle n \rangle} \{u, v\} \\ &= [d'_n, uvd'_{-n}] - \frac{[2n]}{[n]\langle n \rangle} \{u, v\}. \end{aligned}$$

It follows from (3.15) that

$$(3.17) \quad [ud'_n, vd'_{-n}] = [2n]uvd'_0 - \frac{[2n]}{[n]\langle n \rangle} \{u, v\}$$

Similarly, after using quantum Jacobi identity for $(d'_0, ud'_{\pm n}, vd'_{\pm n})$, we obtain

$$(3.18) \quad \langle n \rangle [[d'_0, ud'_{\pm n}], vd'_{\pm n}] + [[ud'_{\pm n}, vd'_{\pm n}], d'_0] + \langle n \rangle [[vd'_{\pm n}, d'_0], ud'_{\pm n}] = 0.$$

Since $\pi([ud'_{\pm n}, vd'_{\pm n}]) = [\pi(ud'_{\pm n}), \pi(vd'_{\pm n})] = [ud_{\pm n}, vd_{\pm n}] = uv[d_{\pm n}, d_{\pm n}] = 0$, $[ud'_{\pm n}, vd'_{\pm n}] \in \mathcal{C}$. It follows from (3.18) that

$$(3.19) \quad [[d'_0, ud'_{\pm n}], vd'_{\pm n}] = [[d'_0, vd'_{\pm n}], ud'_{\pm n}].$$

We have, by (3.10) and (3.19)

$$\begin{aligned} [ud'_{\pm n}, vd'_{\pm n}] &= \mp \frac{1}{[n]} [[d'_0, ud'_{\pm n}], vd'_{\pm n}] \\ &= \mp \frac{1}{[n]} [[d'_0, vd'_{\pm n}], ud'_{\pm n}] \\ &= \mp \frac{1}{[n]} [\mp [n]vd'_{\pm n}, ud'_{\pm n}] \\ &= [vd'_{\pm n}, ud'_{\pm n}] = -[ud'_{\pm n}, vd'_{\pm n}], \end{aligned}$$

i.e.,

$$(3.20) \quad [ud'_{\pm n}, vd'_{\pm n}] = 0.$$

Finally, applying the quantum Jacobi identity to (ud'_0, vd'_n, wd'_{-n}) gives that

$$\{u, vw\} + \{v, wu\} + \{w, uv\} = 0.$$

By the Lemma (2.16) in [3],

$$(3.21) \quad \{u, v\} = -Res \left(u \frac{dv}{dt} \right) \{t, t^{-1}\}, \quad u, v \in \mathbb{C}[t, t^{-1}].$$

Now the Theorem 2.4 follows from (3.9), (3.13), (3.17), (3.20) and (3.21) ■

REFERENCES

1. G.M.Benkart and R.V.Moody, *Derivations, central extensions and affine Lie algebras*, Algebras, Groups and Geometries **3** (1986), 456–492.
2. I.B.Frenkel, J.Lepowsky and A.Meurman, “A vertex operator algebras and the Monster,” Academic Press, New York, 1988.
3. H.Garland, *The arithmetic theory of loop groups*, Inst. Hautes Etudes Sci. Publ. Math. **52** (1980), 5–136.
4. V.G.Kac, “Infinite dimensional Lie algebras,” Cambridge University Press.
5. V.G.Kac, *Highest weight representations of infinite dimensional Lie algebras*, Proc. ICM. Helsinki (1978).
6. Y.I.Manin, “Quantum groups and non-commutative geometry,” CRM-1561, Montreal, 1988.
7. A.Meurman and A.Rocha-Caridi, *Highest weight representations of the Neveu-Schwarz and Ramond algebras*, Commun. Math. Phys. **107** (1986), 263–294.
8. R.V.Moody and A.Pianzola, “Lie algebras with triangular decomposition,” preprint.

CHAPTER 2. THE CHARACTERIZATIONS OF THE QUANTUM WITT ALGEBRA

§2.0. Basic definitions and main theorem

In [1] and [3], two different characterizations of the Witt algebra were found by I. Kaplansky and M.L. Tomber, respectively. In this chapter, we prove that the quantizations of the two characterizations hold for the quantum Witt algebra.

The following notations will be used throughout the paper:

- $\mathbb{Z} :=$ the set of all integers.
- $\mathbb{C} :=$ the set of all complex numbers.
- $[m] := \frac{q^m - q^{-m}}{q - q^{-1}}$, where $m \in \mathbb{Z}$, $q \in \mathbb{C} \setminus \{0\}$ and q is not a root of unity.
- $[0]! := 1$ and $[m]! := [m][m-1] \dots [2][1]$ for $m \in \mathbb{Z}$ and $m > 0$.
- $\langle m \rangle := \frac{q^m + q^{-m}}{2}$, where $m \in \mathbb{Z}$.
- For any $\mathbb{Z}$ -graded vector space $\oplus_{n \in \mathbb{Z}} V_n$, $J \in \text{Hom}(V, V)$ is defined by $J(v_n) := q^n v_n$ for $n \in \mathbb{Z}$ and $v_n \in V_n$.
- $\sigma := \frac{J + J^{-1}}{2}$.
- Let $A = \oplus_{n \in \mathbb{Z}} A_n$ be a $\mathbb{Z}$ -graded algebra over $\mathbb{C}$ with multiplication denoted by xy for $x, y \in A$. As usual, we define the $\mathbb{Z}$ -graded algebras $(A^-, [,])$ and $(A^+, \circ)$ by

$$A^\pm := \oplus_{n \in \mathbb{Z}} (A^\pm)_n, \quad \text{where } (A^\pm)_n := A_n,$$

$$\begin{aligned} [x, y] &:= xy - yx, \\ x \circ y &:= \frac{xy + yx}{2}, \end{aligned}$$

where $x, y \in A$. Let

$$\begin{aligned} (x, y, z)_q &:= (xy)\sigma(z) - \sigma(x)(yz), \\ J_q(x, y, z) &:= (xy)\sigma(z) + (yz)\sigma(z) + (zx)\sigma(y). \end{aligned}$$

where $x, y, z \in A$.

Definition Let A be a $\mathbb{Z}$ -graded algebra over $\mathbb{C}$ with multiplication denoted by xy .

- (1). A is called a **quantum flexible algebra** if $(x, y, x)_q = 0$ for $x, y \in A$.

(2). A is called a **quantum Lie algebra** if $xy = -yx$ and the following **quantum Jacobi identity** holds:

$$J_q(x, y, z) = 0 \quad \text{for } x, y \text{ and } z \in A.$$

One example of quantum Lie algebras is the **quantum Witt algebra** W_q defined by

$$\begin{aligned} W_q &:= \oplus_{n \in \mathbb{Z}} (W_q)_n, & \text{where } (W_q)_n &:= \mathbb{C}d_n, \\ d_m d_n &:= [m - n]d_{m+n}, & \text{where } m, n &\in \mathbb{Z}. \end{aligned}$$

Another interesting example of a quantum Lie algebra is the **quantum Virasoro algebra** $V_q = \oplus_{n \in \mathbb{Z}} (V_q)_n$ defined by

$$(V_q)_n := \begin{cases} \mathbb{C}d_0 \oplus \mathbb{C}c, & \text{if } n = 0, \\ \mathbb{C}d_n, & \text{if } n \neq 0, \end{cases}$$

$$\begin{aligned} d_m d_n &:= [m - n]d_{m+n} + \frac{[m - 1][m][m + 1]}{[2][3]} \delta_{m+n, 0} c, \\ cd_m &:= d_m c := 0, \end{aligned}$$

where $m, n \in \mathbb{Z}$.

Remark. If we define a family of linear operators d_n on the Laurent polynomial ring $\mathbb{C}[t, t^{-1}]$ by

$$d_n := -t^{n+1} \left(\frac{d}{dt} \right)_q, \quad \left(\frac{d}{dt} \right)_q : t^m \mapsto [m]t^{m-1}$$

for all $m, n \in \mathbb{Z}$, then the $\mathbb{Z}$ -graded vector space $\oplus_{n \in \mathbb{Z}} \mathbb{C}d_n$ over $\mathbb{C}$ becomes the quantum Witt algebra under the following multiplication:

$$d_m \cdot d_n := Jd_m J^{-1} d_n J - Jd_n J^{-1} d_m J$$

Now we can state the main theorem.

THEOREM. Let $A = \oplus_{n \in \mathbb{Z}} A_n$ be a $\mathbb{Z}$ -graded algebra over $\mathbb{C}$. Then the following are equivalent:

- (1) A is the quantum Witt algebra.
- (2) A is a quantum flexible algebra and $(A^-, [,]) is the quantum Witt algebra.$
- (3) A is a quantum Lie algebra, $\dim(A_n) \leq 1$ for all $n \in \mathbb{Z}$ and

$$(A_0 A_1) \neq 0, \quad (A_1 A_{-1}) \neq 0, \quad (A_2 A_{-1}) \neq 0, \quad (A_{-2} A_1) \neq 0.$$

Because it is clear that $(1) \Rightarrow (2)$ and $(1) \Rightarrow (3)$, we only need to prove that $(2) \Rightarrow (1)$ and $(3) \Rightarrow (1)$.

§2.1. The proof of the theorem: $(2) \Rightarrow (1)$

Using the same argument as in [2], we can prove:

LEMMA. Let A be a $\mathbb{Z}$ -graded algebra and $(a), (b), (c), (a'), (b')$ denote the following statements:

(a) A is quantum flexible.

(b) $(x, y, z)_q + (z, y, x)_q = 0$ for $x, y, z \in A$.

(c) $[\sigma(x), y \circ z] = [x, y] \circ \sigma(z) + \sigma(y) \circ [x, z]$ for $x, y, z \in A$.

(a') A is quantum flexible and $(A^-, [,]) is a quantum Lie algebra.$

(b') $[\sigma(x), yz] = [x, y]\sigma(z) + \sigma(y)[x, z]$ for $x, y, z \in A$.

Then $(a) \Leftrightarrow (b) \Leftrightarrow (c)$ and $(a') \Leftrightarrow (b')$.

PROOF: $(a) \Rightarrow (b)$: Since A is quantum flexible, we have

$$((x+z)y)\sigma(x+z) = \sigma(x+z)(y(x+z)), \quad \text{where } x, y, z \in A,$$

i.e.

$$\begin{aligned} & (xy)\sigma(x) + (xy)\sigma(z) + (zy)\sigma(x) + (zy)\sigma(z) \\ &= \sigma(x)(yx) + \sigma(x)(yz) + \sigma(z)(yx) + \sigma(z)(yz), \end{aligned}$$

or

$$(2.1) \quad ((xy)\sigma(z) - \sigma(x)(yz)) + ((zy)\sigma(x) - \sigma(z)(yx)) = 0,$$

which is

$$(x, y, z)_q + (z, y, x)_q = 0, \quad \text{where } x, y, z \in A.$$

$(b) \Rightarrow (c)$: It follows from (b) that

$$(x, y, z)_q + (z, y, x)_q + (x, z, y)_q + (y, z, x)_q = (y, x, z)_q + (z, x, y)_q,$$

where $x, y, z \in A$, or

$$\begin{aligned} & (xy)\sigma(z) - \sigma(x)(yz) + (zy)\sigma(x) - \sigma(z)(yx) \\ &+ (xz)\sigma(y) - \sigma(x)(zy) + (yz)\sigma(x) - \sigma(y)(zx) \\ &= (yx)\sigma(z) - \sigma(y)(xz) + (zx)\sigma(y) - \sigma(z)(xy). \end{aligned}$$

This implies that

$$(2.2) \quad [\sigma(x), y \circ z] = [x, y] \circ \sigma(z) + \sigma(y) \circ [x, z] \quad \text{for } x, y, z \in A.$$

(c) $\Rightarrow$ (a): Let $z = x$ in (2.2), we get

$$[\sigma(x), y \circ x] = [x, y] \circ \sigma(x),$$

or

$$\begin{aligned} & \sigma(x)(yx) + \sigma(x)(xy) - (yx)\sigma(x) - (xy)\sigma(x) \\ &= (xy)\sigma(x) - (yx)\sigma(x) + \sigma(x)(xy) - \sigma(x)(yx). \end{aligned}$$

It follows that

$$\sigma(x)(yx) = (xy)\sigma(x) \quad \text{for } x, y \in A.$$

(a') $\Rightarrow$ (b'): Let $x, y, z \in A$. On one hand, by (a) $\Rightarrow$ (c), we have

$$(2.3) \quad [\sigma(x), y \circ z] = [x, y] \circ \sigma(z) + \sigma(y) \circ [x, z].$$

On the other hand, since $(A^-, [,])$ is a quantum Lie algebra, we also have

$$[\sigma(x), [y, z]] + [\sigma(y), [z, x]] + [\sigma(z), [x, y]] = 0,$$

or

$$(2.4) \quad [\sigma(x), \frac{1}{2}[y, z]] = \frac{1}{2}[[x, y]\sigma(z)] + \frac{1}{2}[\sigma(y), [x, z]].$$

It follows from (2.3)+(2.4) that

$$\begin{aligned} [\sigma(x), yz] &= [\sigma(x), y \circ z + \frac{1}{2}[y, z]] = [\sigma(x), y \circ z] + [\sigma(x), \frac{1}{2}[y, z]] \\ &= ([x, y] \circ \sigma(z) + \frac{1}{2}[[x, y], \sigma(z)]) + (\sigma(y) \circ [x, z] + \frac{1}{2}[\sigma(y), [x, z]]) \\ &= [x, y]\sigma(z) + \sigma(y)[x, z]. \end{aligned}$$

(b') $\Rightarrow$ (a'): Let $x = z$ in (b'), we get

$$[\sigma(x), yx] = [x, y]\sigma(x),$$

or

$$\sigma(x)(yx) - (yx)\sigma(x) = (xy)\sigma(x) - (yx)\sigma(x),$$

i.e.,

$$\sigma(x)(yx) = (xy)\sigma(x) \quad \text{for } x, y \in A.$$

This proves that A is a quantum flexible algebra. Therefore, (c) holds for the algebra A .

Next, we prove that $(A^-, [,])$ is a quantum Lie algebra. For $x, y, z \in A$, we have, by (c) and (b')

$$\begin{aligned} \frac{1}{2}[\sigma(x), [y, z]] &= [\sigma(x), \frac{1}{2}[y, z]] \\ &= [\sigma(x), yz - y \circ z] = [\sigma(x), yz] - [\sigma(x), y \circ z] \\ &= ([x, y]\sigma(z) + \sigma(y)[x, z]) - ([x, y] \circ \sigma(z) + \sigma(y) \circ [x, z]) \\ &= ([x, y]\sigma(z) - [x, y] \circ \sigma(z)) + (\sigma(y)[x, z] - \sigma(y) \circ [x, z]) \\ &= \frac{1}{2}[[x, y], \sigma(z)] + \frac{1}{2}[\sigma(y), [x, z]]. \end{aligned}$$

This implies that

$$[[x, y], \sigma(z)] + [[y, z], \sigma(x)] + [[z, x], \sigma(y)] = 0,$$

i.e., $(A^-, [,]) is a quantum Lie algebra. ■$

Now we begin to prove: (2) $\Rightarrow$ (1).

By the Lemma above, we know that (c) and (b') hold for the algebra A in the theorem. Because $(A^-, [,]) is the quantum Witt algebra, we can choose a C-basis $\{e_n \mid n \in \mathbb{Z}\}$ of A such that$

$$\begin{aligned} A &= \bigoplus_{n \in \mathbb{Z}} A_n, \quad \text{where } A_n = \mathbb{C}e_n, \\ [e_n, e_m] &= e_n e_m - e_m e_n = [n - m]e_{n+m}, \quad \text{where } m, n \in \mathbb{Z}, \\ e_m e_n &\in \mathbb{C}e_{m+n}. \end{aligned}$$

In particular, $e_0 e_0 = a e_0$ for some $a \in \mathbb{C}$.

Using (b') to the triple e_n, e_0 and e_m , we get

$$(2.5) \quad \langle n \rangle [e_n, e_0 e_m] = \langle m \rangle [n] e_n e_m + [n - m] e_0 e_{n+m} \quad \text{for } n, m \in \mathbb{Z}.$$

Let $m = 0$ in (2.5), then

$$\langle n \rangle a [n] e_n = [n] (e_n e_0 + e_0 e_n).$$

Hence,

$$(2.6) \quad \langle n \rangle a e_n = e_n e_0 + e_0 e_n \quad \text{for } n \in \mathbb{Z} \setminus \{0\}.$$

On the other hand,

$$(2.7) \quad [n]e_n = e_n e_0 - e_0 e_n \quad \text{for } n \in \mathbb{Z}.$$

It follows from (2.6) and (2.7) that

$$(2.8) \quad e_n e_0 = \frac{\langle n \rangle a + [n]}{2} e_n \quad \text{for } n \in \mathbb{Z} \setminus \{0\}.$$

$$(2.9) \quad e_0 e_n = \frac{\langle n \rangle a - [n]}{2} e_n \quad \text{for } n \in \mathbb{Z} \setminus \{0\}.$$

Let $m = -n \neq 0$ in (2.5), we get

$$\langle n \rangle [e_n, e_0 e_{-n}] = \langle -n \rangle [n] e_n e_{-n} + [2n] e_0 e_0.$$

By (2.9), the equation above becomes

$$(2.10) \quad \frac{\langle n \rangle^2 a + \langle n \rangle [n] - 2a}{2} [2n] e_0 = \langle n \rangle [n] e_n e_{-n} \quad \text{for } n \in \mathbb{Z} \setminus \{0\}.$$

Computing $\frac{\langle n \rangle^2 a + \langle n \rangle [n] - 2a}{2} [2n][n]e_n$ by using (2.8), (2.10) and (b'), we have

$$\begin{aligned} & \frac{\langle n \rangle^2 a + \langle n \rangle [n] - 2a}{2} [2n][n]e_n \\ &= \frac{\langle n \rangle^2 a + \langle n \rangle [n] - 2a}{2} [2n][e_n, e_0] \\ &= [e_n, \langle n \rangle [n] e_n e_{-n}] = [n][\sigma(e_n), e_n e_{-n}] \\ &= \langle n \rangle [n][2n] e_n e_0 \\ &= \frac{1}{2} \langle n \rangle [n][2n](\langle n \rangle a + [n]) e_n \quad \text{for } n \in \mathbb{Z} \setminus \{0\}. \end{aligned}$$

It follows from $\frac{1}{2}[n][2n] \neq 0$ that $a = 0$. Therefore, we have proved that

$$e_n e_0 = -e_0 e_n = \frac{1}{2}[n]e_n \quad \text{for } n \in \mathbb{Z}.$$

Hence,

$$(2.11) \quad e_n \circ e_0 = 0 \quad \text{for } n \in \mathbb{Z}.$$

Now we prove that

$$e_m \circ e_n = 0 \quad \text{for } m, n \in \mathbb{Z}.$$

In fact, we can assume that $m \neq 0$ by (2.11). (c) in the Lemma implies that

$$[\sigma(e_m), e_0 \circ e_n] = [e_m, e_0] \circ \sigma(e_n) + \sigma(e_0) \circ [e_m, e_n].$$

Using (2.11), this gives us the following equation

$$0 = [m] < n > e_m \circ e_n,$$

or

$$e_m \circ e_n = 0.$$

Finally, we get

$$e_m e_n = \frac{1}{2}[e_m, e_n] + e_m \circ e_n = \frac{1}{2}[e_m, e_n] = \frac{1}{2}[m - n]e_{m+n}.$$

Let $d_n := 2^{1-n}e_n$ for $n \in \mathbb{Z}$, then

$$\begin{aligned} A &= \bigoplus_{n \in \mathbb{Z}} A_n, & \text{where } A_n &= \mathbb{C}d_n, \\ d_m d_n &= [m - n]d_{m+n}, & \text{where } m, n &\in \mathbb{Z}, \end{aligned}$$

i.e., A is the quantum Witt algebra.

§2.2. The proof of the theorem: (3) $\Rightarrow$ (1)

As usual, we can choose $0 \neq d_i \in A_i$, where $i = 0, \pm 1$, such that

$$(3.1) \quad d_i d_j = [i - j]d_{i+j} \quad \text{for } i, j = 0, \pm 1.$$

Since $A_2 A_{-1} \neq 0$, $A_2 \neq 0$. Hence, $A_2 = \mathbb{C}u_2$ where $u_2 \neq 0$. Let $d_{-1}u_2 = td_1$ for some $t \in \mathbb{C}$, then $t \neq 0$. Let $d_2 := -\frac{[3]u_2}{t}$, then

$$(3.2) \quad d_2 \neq 0, \quad d_{-1}d_2 = -[3]d_1.$$

For $x \in A$, we define $ad(x) \in \text{End}(A)$ by

$$(ad(x))(a) := xa \quad \text{for } a \in A.$$

For $k \geq 2$, we define

$$(3.3) \quad d_k := \frac{(-1)^k (ad(d_1))^{k-2}(d_2)}{[k-2]}.$$

It is clear that

$$(3.4) \quad d_1 d_{k-1} = [2 - k] d_k \quad \text{for } k \geq 0,$$

Now we use induction on k to prove the following facts:

$$(3.5) \quad d_k \neq 0, \quad d_0 d_k = -[k] d_k, \quad d_{-1} d_k = -[k + 1] d_{k-1} \quad \text{for } k \geq 2,$$

If $k = 2$, then that $d_2 \neq 0$ and $d_{-1} d_2 = -[3] d_1$ follow from (3.2). Let $d_0 d_2 = a_2 d_2$. It follows from $J_q(d_0, d_{-1}, d_2) = 0$ that $a_2 = -[2]$.

Assume that (3.5) is true for $k - 1$ where $k \geq 3$, so there exist $a_k, b_k \in \mathbb{C}$ such that

$$d_0 d_k = a_k d_k, \quad d_{-1} d_k = b_k d_{k-1}.$$

Using (3.4), the induction assumption and $J_q(d_0, d_1, d_{k-1}) = 0$, we have

$$\begin{aligned} 0 &= d_0(d_1 d_{k-1}) + \langle 1 \rangle d_1(d_{k-1} d_0) + \langle k - 1 \rangle d_{k-1}(d_0 d_1) \\ &= [2 - k] d_0 d_k + \langle 1 \rangle [k - 1] d_1 d_{k-1} - \langle k - 1 \rangle d_{k-1} d_1 \\ &= [2 - k] a_k d_k + \langle 1 \rangle [k - 1][2 - k] d_k - \langle k - 1 \rangle [k - 2] d_k. \end{aligned}$$

Since $k \geq 3$, $[2 - k] \neq 0$. Hence, we get

$$(3.6) \quad a_k d_k = (-\langle 1 \rangle [k - 1] - \langle k - 1 \rangle) d_k.$$

Similarly, it follows from $J_q(d_{-1}, d_1, d_{k-1}) = 0$ that

$$(3.7) \quad \langle 1 \rangle [k - 2] b_k d_{k-1} = (-\langle 1 \rangle [k - 3][k] - \langle k - 1 \rangle [2][k - 1]) d_{k-1}.$$

Since $d_{k-1} \neq 0$, it follows from (3.7) that

$$\begin{aligned} \langle 1 \rangle [k - 2] b_k &= -\langle 1 \rangle [k - 3][k] - \langle k - 1 \rangle [2][k - 1] \\ &= -\langle 1 \rangle [k - 2][k + 1]. \end{aligned}$$

So $b_k = -[k + 1] \neq 0$. Hence,

$$d_{-1} d_k = -[k + 1] d_{k-1} \neq 0 \implies d_k \neq 0.$$

Going back to (3.6), we have

$$a_k = -\langle 1 \rangle [k - 1] - \langle k - 1 \rangle = -[k].$$

Therefore, (3.5) holds.

By (3.1), (3.4) and (3.5), we have proved that

$$(3.8) \quad d_i d_k = [i - k] d_{i+k} \quad \text{for } i = 0, \pm 1 \text{ and } k \geq -1.$$

Our next step is to prove that (3.8) is also true for $i = 2$, i.e.

$$(3.9) \quad d_2 d_k = [2 - k] d_{2+k} \quad \text{for } k \geq -1.$$

Let $d_2 d_k = c_k d_{2+k}$ where $c_k \in \mathbb{C}$ and $k \geq -1$. Using (3.8) and $J_q(d_2, d_{-1}, d_k) = 0$, we have

$$\begin{aligned} -\langle 1 \rangle [k + 3] c_k d_{k+1} &= \langle 1 \rangle c_k d_{-1} d_{k+2} = \langle 1 \rangle d_{-1} (d_2 d_k) \\ &= \langle 2 \rangle d_2 (d_{-1} d_k) + \langle k \rangle d_k (d_2 d_{-1}) \\ &= -\langle 2 \rangle [k + 1] d_2 d_{k-1} + \langle k \rangle [3] d_k d_1 \\ &= -\langle 2 \rangle [k + 1] c_{k-1} d_{k+1} + \langle k \rangle [3] [k - 1] d_{k+1}, \end{aligned}$$

i.e.,

$$-\langle 1 \rangle [k + 3] c_k = -\langle 2 \rangle [k + 1] c_{k-1} + \langle k \rangle [3] [k - 1].$$

By (3.8), $c_{-1} = [3]$. Assume that $c_{k-1} = [3 - k]$, then

$$\begin{aligned} -\langle 1 \rangle [k + 3] c_k &= -\langle 2 \rangle [k + 1] [3 - k] + \langle k \rangle [3] [k - 1] \\ &= -\langle 1 \rangle [k + 3] [2 - k]. \end{aligned}$$

So $c_k = [2 - k]$. By induction, (3.9) is true.

Now we can prove that

$$(3.10) \quad d_m d_n = [m - n] d_{m+n} \quad \text{for } m, n \geq -1.$$

Because of (3.8), we can assume that $2 \leq m < n$. We use induction on m . (3.9) tells us that (3.10) is true for $m = 2$. Assume that $m \geq 3$ and (3.10) is true for $m - 1$. From $J_q(d_1, d_{m-1}, d_n) = 0$, we get

$$\langle n \rangle [2 - m] d_m d_n = \langle n \rangle [2 - m] [m - n] d_{m+n}.$$

Since $m \geq 3$, $[2 - m] \neq 0$. It follows that $d_m d_n = [m - n] d_{m+n}$.

Therefore, we have proved that

• **Fact 1.** The subalgebra $\oplus_{n \geq -1} A_n$ of A has a $\mathbb{C}$ -basis

$$\{ d_n \mid n \geq -1, d_n \in A_n \}$$

such that

$$A_n = \mathbb{C}d_n \neq 0, \quad d_m d_n = [m - n]d_{m+n} \quad \text{for } m, n \geq -1.$$

Similarly, if we choose $0 \neq d_{-2} \in A_{-2}$ such that $d_1 d_{-2} = [3]d_1$ and define

$$d_k := \frac{(ad(d_{-1}))^{-k-2}(d_{-2})}{[-k-2]!} \quad \text{for } k \leq -2.$$

Then we can prove the following:

• *Fact 2.* The subalgebra $\oplus_{n \leq 1} A_n$ of A has a $\mathbb{C}$ -basis

$$\{d_n \mid n \leq 1, d_n \in A_n\}$$

such that

$$A_n = \mathbb{C}d_n \neq 0, \quad d_m d_n = [m - n]d_{m+n} \quad \text{for } m, n \leq 1.$$

Furthermore, we have

$$(3.11) \quad d_2 d_{-2} = [4]d_0.$$

In fact, let $d_2 d_{-2} = r d_0$, where $r \in \mathbb{C}$, then

$$(3.12) \quad J_q(d_1, d_{-2}, d_3) = 0 \implies d_4 d_{-2} = [6]d_2.$$

$$(3.13) \quad J_q(d_1, d_{-2}, d_4) = 0 \quad \text{and} \quad (3.12) \implies d_5 d_{-2} = [7]d_3.$$

$$(3.14) \quad J_q(d_1, d_{-2}, d_2) = 0 \implies \langle 2 \rangle d_3 d_{-2} = (\langle 2 \rangle [3]^2 - \langle 1 \rangle r) d_1.$$

Using $J_q(d_2, d_3, d_{-2}) = 0$, (3.13) and (3.14), we get

$$(\langle 1 \rangle + \langle 3 \rangle [3])r = \langle 2 \rangle ([7] + [3]^2),$$

or $\langle 2 \rangle [4]r = \langle 2 \rangle [4]^2$, i.e., $r = [4]$.

In order to complete the proof, we have to prove that

$$(3.15) \quad d_m d_{-n} = [m + n]d_{m-n} \quad \text{for } m, n \in \mathbb{Z}.$$

By fact 1 and fact 2, we can assume that $m \geq 2$ and $n \geq 2$. Now we use induction on m to prove (3.15).

• $m = 2$, in which case, we have to prove

$$(3.16) \quad d_2 d_{-n} = [2 + n] d_{2-n} \quad \text{for } n \geq 2.$$

Let $d_2 d_{-n} = g_n d_{2-n}$, then $g_2 = [4]$ by (3.11). Assuming that $g_n = [2 + n]$, it follows from $J_q(d_{-1}, d_2, d_{-n}) = 0$ that

$$\begin{aligned} < 2 > [n - 1] g_{n+1} = < 1 > [n - 3] g_n + < n > [3][n + 1] \\ &= < 2 > [n - 1][n + 3]. \end{aligned}$$

This implies that $g_{n+1} = [n + 3]$. Hence, we have proved (3.16).

• Assume that (3.15) holds for $2 \leq m \leq k - 1$, then we will prove that (3.15) also holds for $m = k$. In fact, using $J_q(d_1, d_{k-1}, d_{-n}) = 0$, we have

$$< n > [2 - k] d_k d_{-n} = < n > [2 - k][k + n] d_{k-n}.$$

It follows from $< n > [2 - k] \neq 0$ that $d_k d_{-n} = [k + n] d_{k-n}$.

Therefore, we have proved that (3.15). In other words, A is the quantum Witt algebra.

Although the q -analogue of the Theorem 1 in [1] holds for the quantum Witt algebra, the q -analogue of the Theorem 2 in [1] does not hold for the q -analogue of the enveloping algebra of the Witt algebra. In fact, q -deformations of the module of the tensor fields over the Witt algebra are not unique up to isomorphism. We are going to discuss them in chapter 3.

REFERENCES

1. I. Kaplansky, *The Virasoro algebra*, Commu. Math. Phys. **86** (1982), 49-54.
2. H.C. Myung, "Malcev-admissible algebras," Progress in mathematics, Birhauser Boston, Inc., 1986.
3. M.L. Tomber, *Jacobson-Witt algebras and Lie-admissible algebras*, Hadronic J. **4** (1981), 183-197.

CHAPTER 3. QUANTIZATIONS OF THE MODULE OF TENSOR FIELDS OVER THE WITT ALGEBRA

§3.0. Introduction

The representation $V_{\alpha\beta}$ of the Witt algebra on the space of “the tensor fields” of the form $P(z)z^\alpha(dz)^\beta$ is usually called the module of tensor fields over the Witt algebra. Here α and β are complex numbers and $P(z)$ is an arbitrary polynomial in z and z^{-1} . The module $V_{\alpha\beta}$ of the Witt algebra plays a very important role in the representation theory of the Virasoro algebra. In 1982, I.Kaplansky proved in [3] that if $\oplus_{n \in \mathbb{Z}} \mathbb{C}v_n$ is a $\mathbb{Z}$ -graded module of the Witt algebra $W = \oplus_{n \in \mathbb{Z}} \mathbb{C}d_n$ and $d_{\pm 1}$ are injective operators on V , then V is isomorphic to the module $V_{\alpha\beta}$ of tensor fields for some $\alpha, \beta \in \mathbb{C}$. We call this result Kaplansky’s Theorem. The main purpose of this chapter is to prove a q -analogue of Kaplansky’s Theorem.

Throughout this chapter, we assume that

- All vector spaces are the vector spaces over the complex number field $\mathbb{C}$.
- $\mathbb{C}^* := \{x \in \mathbb{C} \mid x \neq 0\}$.
- q is a complex number satisfying $q^2 \neq 0, 1$.
- $\ell n(z)$ is the principal value of the logarithm.
- $q^\alpha := e^{\alpha \ell n(q)}$ for $\alpha \in \mathbb{C}$.
- $[\alpha] := \frac{q^\alpha - q^{-\alpha}}{q - q^{-1}}$ for $\alpha \in \mathbb{C}$.

In section 1, after defining q -analogue $U(W_q)$ of the enveloping algebra of the Witt algebra, we will construct two kinds of $U(W_q)$ -modules $A(\lambda, \alpha, \beta)$ and $B(\lambda, \alpha, \beta)$ by using a version of the operations over $\mathbb{Z}$ -graded modules of the Witt algebra introduced by B.L.Feigin and D.B.Fuchs in [1], where $(\lambda, \alpha, \beta) \in \mathbb{C}^* \times \mathbb{C} \times \mathbb{C}$. Both $A(\lambda, \alpha, \beta)$ and $B(\lambda, \alpha, \beta)$ become the module of tensor fields over the Witt algebra when $q \rightarrow 1$. In section 2, we will find the necessary and sufficient conditions for $X(\lambda, \alpha, \beta) \simeq Y(\lambda', \alpha', \beta')$ (where $X, Y \in \{A, B\}$) and study the reducibility and unitarity of $X(\lambda, \alpha, \beta)$. In section 3, we will prove a q -analogue of Kanplansky’s theorem.

§3.1. The construction of $U(W_q)$ -modules $X(\lambda, \alpha, \beta)$

Based on Proposition 1.1 in chapter 1, we introduce the following definition:

Definition 1.1 q -analogue $U(W_q)$ of the enveloping algebra of the Witt algebra is defined as the associative algebra with generators $\{J^{\pm 1}, d_n \mid n \in \mathbb{Z}\}$ and the following relations:

$$(1.1) \quad JJ^{-1} = J^{-1}J = 1, \quad Jd_mJ^{-1} = q^m d_m,$$

$$(1.2) \quad q^m d_m d_n J - q^n d_n d_m J = [m - n] d_{m+n},$$

where $m, n \in \mathbb{Z}$.

Definition 1.2 A $U(W_q)$ -module V is called a $\mathbb{Z}$ -graded module if $V = \bigoplus_{n \in \mathbb{Z}} V_n$ and $d_m(v_n) \in V_{m+n}$ for $m, n \in \mathbb{Z}$.

For every $\lambda \in \mathbb{C}^*$, we define an algebra isomorphism $\varphi(\lambda)$ of $U(W_q)$ as follows:

$$\varphi(\lambda) : J^{\pm 1} \mapsto \lambda^{\pm 1} J^{\pm 1}, \quad d_m \mapsto \lambda^{-1} d_m \quad \text{for } m \in \mathbb{Z}.$$

If $V = \bigoplus_{n \in \mathbb{Z}} V_n$ is a $\mathbb{Z}$ -graded $U(W_q)$ -module with $J(v_n) = q^n v_n$ for all $n \in \mathbb{Z}$ and $v_n \in V_n$, then we can construct three more modules from V : **contragradient module** $\bar{V} := \bigoplus_{n \in \mathbb{Z}} (\bar{V})_n$, **adjoint module** $V^* := \bigoplus_{n \in \mathbb{Z}} (V^*)_n$ and **inverted module** $V^\circ := \bigoplus_{n \in \mathbb{Z}} (V^\circ)_n$, where

$$\begin{aligned} (\bar{V})_n &:= \text{Hom}(V_n, \mathbb{C}), & J|(\bar{V})_n &:= q^n \cdot \text{id}; \\ (V^*)_n &:= \text{Hom}(V_{-n}, \mathbb{C}), & J|(V^*)_n &:= q^{-n} \cdot \text{id}; \\ (V^\circ)_n &:= V_{-n}, & J|(V^\circ)_n &:= q^{-n} \cdot \text{id} \end{aligned}$$

and the definitions of the operators d_m on $\bar{V}$, V^* and V° are the same as in [1].

It is easy to check that $\bar{V}$ is a $\mathbb{Z}$ -graded $U(W_q)$ -module and V^* , as well as V° , is a $\mathbb{Z}$ -graded $U(W_{q^{-1}})$ -module. As $U(W_q)$ -modules, $(V^*)^\circ \simeq \bar{V}$.

In particular, if $V = \bigoplus_{k \in \mathbb{Z}} \mathbb{C} v_k$ is a $\mathbb{Z}$ -graded $U(W_q)$ -module with the natural $\mathbb{Z}$ -grading and the following module action on V :

$$(1.3) \quad J(v_k) := q^k v_k, \quad d_n(v_k) := a(q, n, k) v_{n+k}$$

where $n, k \in \mathbb{Z}$ and $a(q, n, k) \in \mathbb{C}$, then we can describe the contragradient module $\bar{V}$, the adjoint module V^* and the inverted module V° as follows:

$$(1.4) \quad \bar{V} = \bigoplus_{k \in \mathbb{Z}} \mathbb{C} v_k, \quad J(v_k) = q^k v_k, \quad d_n(v_k) = a(q, -n, n+k) v_{n+k};$$

$$(1.5) \quad V^* = \bigoplus_{k \in \mathbb{Z}} \mathbb{C} v_k, \quad J(v_k) = q^{-k} v_k, \quad d_n(v_k) = -a(q, n, -n-k) v_{n+k};$$

$$(1.6) \quad V^\circ = \oplus_{k \in \mathbb{Z}} \mathbb{C} v_k, \quad J(v_k) = q^{-k} v_k, \quad d_n(v_k) = -a(q, -n, -k) v_{n+k}.$$

For $\alpha, \beta \in \mathbb{C}$, set

$$(1.7) \quad a(q, n, k) := -([k + \alpha]q^\alpha + [n + 1][\beta]q^{n+k}),$$

where $n, k \in \mathbb{Z}$. Then (1.3) and (1.7) define a $U(W_q)$ -module action on $V(\alpha, \beta) := \oplus_{k \in \mathbb{Z}} \mathbb{C} v_k$.

Let us check (1.2), i.e.

$$(1.8) \quad q^m d_m d_n(v_k) - q^n d_n d_m(v_k) = [m - n] d_{m+n} J^{-1}(v_k) \quad \text{for } m, n, k \in \mathbb{Z}.$$

Let $q^m d_m d_n(v_k) = A_{m,n,k} v_{m+n+k}$, then $q^n d_n d_m(v_k) = A_{n,m,k} v_{m+n+k}$. By (1.7), we have

$$\begin{aligned} A_{m,n,k} &= q^{m+2\alpha}[k + \alpha][n + k + \alpha] + q^{2m+n+k+\alpha}[k + \alpha][m + 1][\beta] \\ &\quad + q^{m+n+k+\alpha}[n + 1][\beta][n + k + \alpha] \\ &\quad + q^{2m+2n+2k}[n + 1][m + 1][\beta]^2 \end{aligned}$$

It follows that

$$\begin{aligned} q^m d_m d_n(v_k) - q^n d_n d_m(v_k) &= (A_{m,n,k} - A_{n,m,k}) v_{m+n+k} \\ &= ([k + \alpha](q^{m+2\alpha}[n + k + \alpha] - q^{n+2\alpha}[m + k + \alpha]) \\ &\quad + q^{m+n+k}[\beta]((q^{m+\alpha}[k + \alpha][m + 1] + q^\alpha[n + 1][n + k + \alpha]) \\ &\quad - (q^{n+\alpha}[k + \alpha][n + 1] + q^\alpha[m + 1][m + k + \alpha])) v_{m+n+k} \\ &= ([k + \alpha](-q^{-k} q^\alpha [m - n]) \\ &\quad + q^{m+n+k}[\beta](-q^{-k} [m - n][m + n + 1])) v_{m+n+k} \\ &= -[m - n]([k + \alpha]q^\alpha + [m + n + 1][\beta]q^{m+n+k}) q^{-k} v_{m+n+k} \\ &= [m - n] d_{m+n} J^{-1}(v_k), \end{aligned}$$

so (1.8) is true.

By the discussion above, we can construct the contragredient module $\bar{V}(\alpha, \beta)$ from (1.4). By (1.7),

$$\begin{aligned} -a(q, -n, n + k) &= [n + k + \alpha]a^\alpha + [-n + 1][\beta]q^k \\ &= ([n + k]q^\alpha + [\alpha]q^{-n-k})q^\alpha + [\beta]\frac{q^{1-n} - q^{n-1}}{q - q^{-1}}q^k \\ &= \frac{q^{n+k} - q^{-n-k}}{q - q^{-1}}q^{2\alpha} + \frac{q^{2\alpha} - 1}{q - q^{-1}}q^{-n-k} + \frac{[\beta]q}{q - q^{-1}}q^{-n+k} - \frac{[\beta]q^{-1}}{q - q^{-1}}q^{n+k} \\ &= \frac{q^{-n+k}}{q - q^{-1}}((q^{2n} - q^{-2k})q^{2\alpha} + (q^{2\alpha} - 1)q^{-2k} + [\beta]q - [\beta]q^{-1}q^{2n}) \\ &= \frac{q^{-n+k}}{q - q^{-1}}(-q^{-2k} + (q^{2\alpha} - [\beta]q^{-1})q^{2n} + [\beta]q). \end{aligned}$$

Hence, the contragradient module $\bar{V}(\alpha, \beta)$ can be described as

$$(1.9) \quad \begin{aligned} \bar{V}(\alpha, \beta) &:= \oplus_{k \in \mathbb{Z}} \mathbb{C} u_k, & J(u_k) &:= q^k u_k, \\ d_n(u_k) &:= -([n+k+\alpha]q^\alpha + [1-n][\beta]q^k)u_{n+k} \\ &= -\frac{q^{-n+k}}{q-q^{-1}}(-q^{-2k} + (q^{2\alpha} - [\beta]q^{-1})q^{2n} + [\beta]q)u_{n+k}. \end{aligned}$$

If we replace q by q^{-1} in (1.5), then we get a $\mathbb{Z}$ -graded $U(W_q)$ -module $V(\alpha, \beta)^{(1)}$:

$$(1.10) \quad \begin{aligned} V(\alpha, \beta)^{(1)} &:= \oplus_{k \in \mathbb{Z}} \mathbb{C} u_k, & J(u_k) &:= q^k u_k, \\ d_n(u_k) &:= -a(q^{-1}, n, -n-k)u_{n+k} \\ &= ([-n-k+\alpha]q^{-n} + [n+1][\beta]q^k)u_{n+k} \\ &= -\frac{q^{-n+k}}{q-q^{-1}}(-q^{-2k} + (q^{-2\alpha} - q[\beta])q^{2n} + q^{-1}[\beta])u_{n+k}. \end{aligned}$$

Similarly, if we replace q by q^{-1} in (1.6), then, by (1.7)

$$\begin{aligned} a(q^{-1}, -n, -k) &= -([-k+\alpha]q^{-\alpha} + [-n+1][\beta]q^{n+k}) \\ &= [k-\alpha]q^{-\alpha} + [n-1][\beta]q^{n+k} \\ &= \frac{q^k}{q-q^{-1}}(-q^{-2k} + q^{-1}[\beta]q^{2n} + (q^{-2\alpha} - q[\beta])). \end{aligned}$$

Choose $\alpha', \beta' \in \mathbb{C}$ such that

$$q[\beta'] = q^{-2\alpha} - q[\beta] \quad \text{and} \quad q^{2\alpha'} - q^{-1}[\beta'] = q^{-1}[\beta],$$

then we get a $\mathbb{Z}$ -graded $U(W_q)$ -module $V(\alpha, \beta)^{(2)}$ as follows :

$$(1.11) \quad \begin{aligned} V(\alpha, \beta)^{(2)} &:= \oplus_{k \in \mathbb{Z}} \mathbb{C} v_k, & J(v_k) &:= q^k v_k, \\ d_n(v_k) &:= -([n+k+\alpha]q^{n+\alpha} + [1-n][\beta]q^{n+k})v_{n+k} \\ &= -\frac{q^k}{q-q^{-1}}(-q^{-2k} + (q^{2\alpha} - q^{-1}[\beta])q^{2n} + q[\beta])v_{n+k}. \end{aligned}$$

Finally, let us rewrite the $\mathbb{Z}$ -graded $U(W_q)$ -module as:

$$(1.12) \quad \begin{aligned} V(\alpha, \beta) &:= \oplus_{k \in \mathbb{Z}} \mathbb{C} v_k, & J(v_k) &:= q^k v_k, \\ d_n(v_k) &:= -([k+\alpha]q^\alpha + [1+n][\beta]q^{n+k})v_{n+k} \\ &= -\frac{q^k}{q-q^{-1}}(-q^{-2k} + q[\beta]q^{2n} + (q^{2\alpha} - q^{-1}[\beta]))v_{n+k}. \end{aligned}$$

Let

$$v_k := q^{-k} u_k \quad \text{for } k \in \mathbb{Z}.$$

By (1.9) and (1.11)

$$(1.13) \quad \bar{V}(\alpha, \beta) = \oplus_{k \in \mathbb{Z}} \mathbb{C} v_k = V(\alpha, \beta)^{(2)}.$$

If we choose $\alpha', \beta' \in \mathbb{C}$ such that

$$q[\beta'] = q^{-2\alpha} - q[\beta] \quad \text{and} \quad q^{2\alpha'} - q^{-1}[\beta'] = q^{-1}[\beta],$$

then, by (1.10) and (1.12)

$$(1.14) \quad V(\alpha, \beta)^{(1)} = \oplus_{k \in \mathbb{Z}} \mathbb{C} v_k = V(\alpha', \beta').$$

(1.13) and (1.14) implies that the contraction which produces the module $\bar{V}(\alpha, \beta)$ (resp. $V(\alpha, \beta)^{(1)}$) does not take us out of the class of the $\mathbb{Z}$ -graded $U(W_q)$ -module $V(\alpha, \beta)^{(2)}$ (resp. $V(\alpha, \beta)$).

Hence, for any $(\lambda, \alpha, \beta) \in \mathbb{C}^* \times \mathbb{C} \times \mathbb{C}$, we can construct two kinds of $\mathbb{Z}$ -graded $U(W_q)$ -module $A(\lambda, \alpha, \beta)$ and $B(\lambda, \alpha, \beta)$ by using (1.11), (1.12) and $\varphi(\lambda)$ as follows (where $n, k \in \mathbb{Z}$):

$$(1.15) \quad \begin{aligned} A(\lambda, \alpha, \beta) &:= \oplus_{k \in \mathbb{Z}} \mathbb{C} v_k, & J(v_k) &:= \lambda q^k v_k, \\ d_n(v_k) &:= -\lambda^{-1} ([k + \alpha] q^\alpha + [1 + n][\beta] q^{n+k}) v_{n+k} \\ &= -\frac{\lambda^{-1} q^k}{q - q^{-1}} (-q^{-2k} + q[\beta] q^{2n} + (q^{2\alpha} - q^{-1}[\beta])) v_{n+k} \end{aligned}$$

and

$$(1.16) \quad \begin{aligned} B(\lambda, \alpha, \beta) &:= \oplus_{k \in \mathbb{Z}} \mathbb{C} v_k, & J(v_k) &:= \lambda q^k v_k, \\ d_n(v_k) &:= -\lambda^{-1} ([n + k + \alpha] q^{n+\alpha} + [1 - n][\beta] q^{n+k}) v_{n+k} \\ &= -\frac{\lambda^{-1} q^k}{q - q^{-1}} (-q^{-2k} + (q^{2\alpha} - q^{-1}[\beta]) q^{2n} + q[\beta]) v_{n+k} \end{aligned}$$

§3.2. The properties of $U(W_q)$ -modules $X(\lambda, \alpha, \beta)$

In this section, we assume that q is not a root of unity.

Let X be A or B , we define

$$\mathcal{cl}(X) := \{ X(\lambda, \alpha, \beta) \mid (\lambda, \alpha, \beta) \in \mathbb{C}^* \times \mathbb{C} \times \mathbb{C} \}.$$

A $U(W_q)$ -module V is said to be in $\mathcal{cl}(X)$ if $V \simeq X(\lambda, \alpha, \beta)$ (as $U(W_q)$ -modules) for some $X(\lambda, \alpha, \beta) \in \mathcal{cl}(X)$.

PROPOSITION 2.1. *For any fixed $h \in \mathbb{Z}$, $X(\lambda, \alpha, \beta) \simeq X(\lambda q^h, \alpha + h, \beta')$ as $U(W_q)$ -modules, where $\beta' \in \mathbb{C}$ with $[\beta'] = [\beta]q^{2h}$.*

PROOF: Let

$$X(\lambda, \alpha, \beta) = \oplus_{k \in \mathbb{Z}} \mathbb{C} v_k \quad \text{and} \quad X(\lambda q^h, \alpha + h, \beta') = \oplus_{k \in \mathbb{Z}} \mathbb{C} u_k.$$

We define a linear map $\varphi : X(\lambda, \alpha, \beta) \rightarrow X(\lambda q^h, \alpha + h, \beta')$ by

$$\varphi : v_k \mapsto u_{k-h} \quad \text{for } k \in \mathbb{Z}.$$

It is clear that φ is bijective. Now we prove that φ preserves the $U(W_q)$ -module actions.

From

$$\begin{aligned} \varphi J(v_k) &= \varphi(\lambda q^k v_k) = \lambda q^k \varphi(v_k) \\ &= \lambda q^k u_{k-h} = (\lambda q^h)(q^{k-h} u_{k-h}) \\ &= J(u_{k-h}) = J\varphi(v_k), \end{aligned}$$

we get that $\varphi J(v_k) = J\varphi(v_k)$.

Let

$$\beta' := \frac{\ell n([\beta]q^{2h} + \sqrt{[\beta]^2 q^{4h}(q - q^{-1})^2 + 4}) - \ell n(2)}{\ell n q},$$

then $[\beta'] = [\beta]q^{2h}$.

If $X = A$, then, by (1.15)

$$\begin{aligned} \varphi d_n(v_k) &= -\frac{\lambda^{-1} q^k}{q - q^{-1}} (-q^{-2k} + q[\beta]q^{2n} + (q^{2\alpha} - q^{-1}[\beta])) \varphi(v_{n+k}) \\ &= -\frac{\lambda^{-1} q^k}{q - q^{-1}} (-q^{-2k} + q[\beta]q^{2n} + (q^{2\alpha} - q^{-1}[\beta])) u_{n+k-h} \\ &= -\frac{(\lambda q^h)^{-1} q^{k-h}}{q - q^{-1}} (-q^{-2(k-h)} + q[\beta']q^{2n} + (q^{2(\alpha+h)} - q^{-1}[\beta'])) u_{n+k-h} \\ &= d_n(u_{k-h}) = d_n \varphi(v_k). \end{aligned}$$

If $X = B$, then, by (1.16)

$$\begin{aligned}
\varphi d_n(v_k) &= -\frac{\lambda^{-1} q^k}{q - q^{-1}} (-q^{-2k} + (q^{2\alpha} - q^{-1}[\beta])q^{2n} + q[\beta])\varphi(v_{n+k}) \\
&= -\frac{\lambda^{-1} q^k}{q - q^{-1}} (-q^{-2k} + (q^{2\alpha} - q^{-1}[\beta])q^{2n} + q[\beta])u_{n+k-h} \\
&= -\frac{(\lambda q^k)^{-1} q^{k-h}}{q - q^{-1}} (-q^{-2(k-h)} + (q^{2(\alpha+h)} - q^{-1}[\beta'])q^{2n} + q[\beta'])u_{n+k-h} \\
&= d_n(u_{k-h}) = d_n\varphi(v_k).
\end{aligned}$$

This proves Proposition 2.1. ■

PROPOSITION 2.2. If $q \in \mathbb{R}$, then $U(W_q)$ has an antilinear anti-involution θ such that $\theta(J^{\pm 1}) := J^{\pm 1}$ and $\theta(d_n) := d_{-n}$ for all $n \in \mathbb{Z}$.

PROOF: Let F be the free associative algebra on the set $\{ \underline{J}^{\pm 1}, \underline{d}_m \mid m \in \mathbb{Z} \}$. Let I be the ideal of F generated by

$$\begin{aligned}
\underline{J} \cdot \underline{J}^{-1} - 1, \quad \underline{J}^{-1} \cdot \underline{J} - 1, \quad \underline{J} \underline{d}_m \underline{J}^{-1} - q^m \underline{d}_m, \\
q^m \underline{d}_m \cdot \underline{d}_n \underline{J} - q^n \underline{d}_n \cdot \underline{d}_m \underline{J} - [m - n] \underline{d}_{m+n},
\end{aligned}$$

where $m, n \in \mathbb{Z}$. Then $U(W_q) := \frac{F}{I}$.

Define an antilinear map $\theta : F \rightarrow U(W_q)$ by

$$\begin{aligned}
\theta(\underline{J}^{\pm 1}) &:= J, \quad \theta(\underline{d}_m) := d_{-m} \\
\theta : a \underline{x}_1 \cdot \underline{x}_2 \dots \underline{x}_t &\mapsto \bar{a} x_t \dots x_2 x_1,
\end{aligned}$$

where $a \in \mathbb{C}$, $x_i \in \{ J^{\pm 1}, d_m \mid m \in \mathbb{Z} \}$, $x_i := \underline{x}_i + I$ and t is a natural number. Then

$$\theta(uv) = \theta(v)\theta(u) \quad \text{for } u, v \in F.$$

If $q \in \mathbb{R}$, then $[m] \in \mathbb{R}$ for $m \in \mathbb{Z}$. So

$$\begin{aligned}
\theta(q^m \underline{d}_m \cdot \underline{d}_n \underline{J} - q^n \underline{d}_n \cdot \underline{d}_m \underline{J}) &= q^m \theta(\underline{J}) \theta(\underline{d}_n) \theta(\underline{d}_m) - q^n \theta(\underline{J}) \theta(\underline{d}_m) \theta(\underline{d}_n) \\
&= q^m (J d_{-n}) d_{-m} - q^n (J d_{-m}) d_{-n} \\
&= q^m (q^{-n} d_{-n} J) d_{-m} - q^n (q^{-m} d_{-m} J) d_{-n} \\
&= q^{m-n} d_{-n} (J d_{-m}) - q^{n-m} d_{-m} (J d_{-n}) \\
&= q^{m-n} d_{-n} (q^{-m} d_{-m} J) - q^{n-m} d_{-m} (q^{-n} d_{-n} J) \\
&= q^{-n} d_{-n} d_{-m} J - q^{-m} d_{-m} d_{-n} J \\
&= [m - n] d_{-(m+n)} = [m - n] \theta(\underline{d}_{m+n}) \\
&= \theta([m - n] \underline{d}_{m+n}),
\end{aligned}$$

or

$$\theta(q^m \underline{d_m} \cdot \underline{d_n} J - q^n \underline{d_n} \cdot \underline{d_m} J - (m-n) \underline{d_{m+n}}) = 0.$$

Similarly, we have

$$\theta(\underline{J} \cdot \underline{J^{-1}} - 1) = \theta(\underline{J^{-1}} \cdot \underline{J} - 1) = \theta(\underline{J} \underline{d_m} \underline{J^{-1}} - q^m \underline{d_m}) = 0.$$

Therefore, θ vanishes on I . This induces an antilinear anti-homomorphism $\theta : U(W_q) \rightarrow U(W_q)$ such that

$$\theta(J_{\pm 1}) = J^{\pm 1} \quad \text{and} \quad \theta(d_n) = d_{-n} \quad \text{for } n \in \mathbb{Z}.$$

It is clear that $\theta^2 = 1$. So, θ is an anlinear anti-involution. ■

Definition 2.1 Let $q \in \mathbb{R}$. A $U(W_q)$ -module V is **unitary** with respect to θ if there is an Hermitian form $\langle \cdot | \cdot \rangle$ on V such that

$$\langle v | v \rangle > 0 \quad \text{for } v \in V \text{ and } v \neq 0,$$

$$(*) \quad \langle x(u) | v \rangle = \langle u | \theta(x)v \rangle \quad \text{for } u, v \in V \text{ and } x \in U(W_q).$$

An Hermitian form $\langle \cdot | \cdot \rangle$ satisfying $(*)$ is called a **contravariant form**.

Let $X(\alpha, \beta) := X(1, \alpha, \beta)$, then the following proposition is clear:

PROPOSITION 2.3. For $(\lambda, \alpha, \beta) \in \mathbb{C}^* \times \mathbb{C} \times \mathbb{C}$, we have

- (1) $X(\lambda, \alpha, \beta)$ is reducible if and only if $X(\alpha, \beta)$ is reducible.
- (2) If $q \in \mathbb{R}$, then $X(\lambda, \alpha, \beta)$ is unitary with respect to θ if and only if $\lambda \in \mathbb{R}$, $\lambda \neq 0$ and $X(\alpha, \beta)$ is unitary with respect to θ . ■

Now we prove

PROPOSITION 2.4. Let $X, Y \in \{A, B\}$. Then

- (1) $X(\lambda, \alpha, \beta) \simeq Y(\lambda_1, \alpha_1, \beta_1) \iff$ there exist some $h \in \mathbb{Z}$ such that $\lambda = \lambda_1 q^h$ and some $\mathbb{Z}$ -grading preserving isomorphism φ such that $\varphi : X(\alpha, \beta) \simeq Y(\alpha', \beta')$, where $\alpha' = \alpha_1 + h$ and $[\beta'] = [\beta_1] q^{2h}$.
- (2) Every submodule of $X(\lambda, \alpha, \beta)$ respects the $\mathbb{Z}$ -grading of $X(\lambda, \alpha, \beta)$.

PROOF: (1). $\Rightarrow$: Let

$$X(\lambda, \alpha, \beta) = \oplus_{k \in \mathbb{Z}} \mathbb{C} v_k, \quad Y(\lambda_1, \alpha_1, \beta_1) = \oplus_{k \in \mathbb{Z}} \mathbb{C} u_k,$$

$$\psi : X(\lambda, \alpha, \beta) \simeq Y(\lambda_1, \alpha_1, \beta_1), \quad \psi(v_k) = a_{j_1} u_{j_1} + \cdots + a_{j_r} u_{j_r},$$

where $a_{j_s} \in \mathbb{C}^*$ and $j_s \neq j_t$ if $s \neq t$. That $\psi J(v_k) = J\psi(v_k)$ gives that $\lambda = \lambda_1 q^{j_s - k}$ for all $1 \leq s \leq r$. Because q is not a root of unity, $r = 1$. It follows that

$$\psi(v_k) = a_{f(k)} u_{f(k)}, \quad \text{where } f(k) \in \mathbb{Z}.$$

Since $q^{f(k)-k} = \frac{\lambda}{\lambda_1}$, $f(k) - k = f(k') - k'$ for all $k, k' \in \mathbb{Z}$. Let $h := f(k) - k$, then

$$\psi(v_k) = a_{k+h} u_{k+h} \quad \text{for } k \in \mathbb{Z}.$$

By Proposition 2.1, $\eta : Y(\lambda_1, \alpha_1, \beta_1) \simeq Y(\lambda_1 q^h, \alpha', \beta')$, where $\alpha' = \alpha_1 + h$ and $[\beta'] = [\beta_1] q^{2h}$. Let $\varphi := \eta\psi$, then φ preserves the $\mathbb{Z}$ -grading and $\varphi : X(\lambda, \alpha, \beta) \simeq Y(\lambda, \alpha', \beta')$. Using the automorphism $\varphi(\lambda)$ of $U(W_q)$, we get that $\varphi : X(\alpha, \beta) \simeq Y(\alpha', \beta')$.

$\Leftarrow$: If $X(\alpha, \beta) \simeq Y(\alpha', \beta')$, then

$$\begin{aligned} X(\lambda, \alpha, \beta) &\simeq Y(\lambda, \alpha', \beta') \\ &= Y(\lambda_1 q^h, \alpha_1 + h, \beta') \simeq Y(\lambda_1, \alpha_1, \beta_1). \end{aligned}$$

(2) Let V be a submodule of $X(\lambda, \alpha, \beta) = \bigoplus_{k \in \mathbb{Z}} \mathbb{C} v_k$.

If $v \in V$, then

$$v = a_{j_1} v_{j_1} + \dots + a_{j_r} v_{j_r},$$

where $a_{j_s} \in \mathbb{C}^*$ and $j_s \neq j_t$ if $1 \leq s \neq t \leq r$.

Applying the operator J to $v, J(v), \dots, J^{r-2}(v)$, respectively, we get a system with respect to $a_{j_1} v_{j_1}, a_{j_2} v_{j_2}, \dots, a_{j_r} v_{j_r}$:

$$\begin{aligned} v &= a_{j_1} v_{j_1} + a_{j_2} v_{j_2} + \dots + a_{j_r} v_{j_r}, \\ \lambda^{-1} J(v) &= q^{j_1} (a_{j_1} v_{j_1}) + q^{j_2} (a_{j_2} v_{j_2}) + \dots + q^{j_r} (a_{j_r} v_{j_r}), \end{aligned}$$

.....

$$\lambda^{-(r-1)} J^{r-1}(v) = (q^{j_1})^{r-1} (a_{j_1} v_{j_1}) + (q^{j_2})^{r-1} (a_{j_2} v_{j_2}) + \dots + (q^{j_r})^{r-1} (a_{j_r} v_{j_r}).$$

The determinant of its coefficients matrix is a Vandermonde determinant with respect to $q^{j_1}, q^{j_2}, \dots, q^{j_r}$. Since $q^{j_s} \neq q^{j_t}$ for $1 \leq s \neq t \leq r$, the determinant is not zero. Thus each of the $a_{j_s} v_{j_s}$ ($s = 1, 2, \dots, r$) can be expressed as a linear combination of $v, \lambda^{-1} J(v), \dots, \lambda^{-(r-1)} J^{r-1}(v)$, i.e., $a_{j_s} v_{j_s} \in V$, or $v_{j_s} \in V$, proving (2). ■

The proposition above tells us that if q is not a root of unity, then in order to study the properties of the $U(W_q)$ -module $X(\lambda, \alpha, \beta)$, it suffices to study the properties of the $U(W_q)$ -module $X(\alpha, \beta)$.

PROPOSITION 2.5. Let $(\alpha, \beta) \in \mathbb{C} \times \mathbb{C}$. Then $B(\alpha, \beta) \in \text{cl}(A) \iff q^{2\alpha+1} + (q^4 - 1)[\beta] \neq 0$.

PROOF: $\Leftarrow$: Since $q^{2\alpha+1} + (q^4 - 1)[\beta] \neq 0$, we can find $(\alpha', \beta') \in \mathbb{C} \times \mathbb{C}$ such that

$$q^{2\alpha'} - q^{-1}[\beta'] = q[\beta] \quad \text{and} \quad q[\beta'] = q^{2\alpha} - q^{-1}[\beta].$$

Hence, $B(\alpha, \beta) = A(\alpha', \beta') \in \text{cl}(A)$ by (1.15) and (1.16).

$\Rightarrow$: If $B(\alpha, \beta) \in \text{cl}(A)$, then, by Proposition 2.4, there exists a $\mathbb{Z}$ -grading preserving isomorphism φ such that

$$\varphi : A(\alpha', \beta') \simeq B(\alpha, \beta) \quad \text{for some } (\alpha', \beta') \in \mathbb{C} \times \mathbb{C}.$$

Set

$$\begin{aligned} A(\alpha', \beta') &= \oplus_{k \in \mathbb{Z}} \mathbb{C} v'_k, & B(\alpha, \beta) &= \oplus_{k \in \mathbb{Z}} \mathbb{C} v_k, \\ \varphi(v'_k) &= a_k v_k, & \text{where } a_k &\in \mathbb{C}, \end{aligned}$$

and

$$(2.1) \quad a_{\alpha\beta} := q^{2\alpha} - q^{-1}[\beta], \quad b_{\alpha,\beta} := a_{\alpha,\beta} - 1, \quad c_\beta := q[\beta] - 1$$

Using (1.15), (1.16) and $\varphi d_n(v'_k) = d_n \varphi(v'_k)$, we have

$$(2.2) \quad (-x + q[\beta']y + a_{\alpha'\beta'})a_{n+k} = (-x + a_{\alpha\beta}y + q[\beta])a_k,$$

where

$$x := q^{-2k} \quad \text{and} \quad y := q^{2n}.$$

It follows from (2.2) that

$$(2.3) \quad (b_{\alpha'\beta'}x + q[\beta']y)a_{n+k} = (c_\beta x + a_{\alpha\beta}y)a_0,$$

$$(2.4) \quad (b_{\alpha'\beta'}x + q[\beta']y)a_k = (c_\beta x + a_{\alpha\beta}y)a_0.$$

Multiplying the both sides of (2.2) by

$$(b_{\alpha'\beta'}x + q[\beta']y)(b_{\alpha'\beta'}x + q[\beta']),$$

we get by using (2.3) and (2.4)

$$\begin{aligned} &(-x + q[\beta']y + a_{\alpha'\beta'})(c_\beta x + a_{\alpha\beta}y)(b_{\alpha'\beta'}x + q[\beta']) \\ &= (-x + a_{\alpha\beta}y + q[\beta])(b_{\alpha'\beta'}x + q[\beta']y)(c_\beta x + a_{\alpha\beta}y). \end{aligned}$$

Comparing the coefficients of x^2y and xy^2 gives us the following identities:

$$(2.5) \quad -a_{\alpha\beta}b_{\alpha'\beta'} + q[\beta']c_{\beta}b_{\alpha'\beta'} = -q[\beta']c_{\beta} + a_{\alpha\beta}b_{\alpha'\beta'}c_{\beta},$$

$$(2.6) \quad [\beta']a_{\alpha\beta}b_{\alpha'\beta'} = [\beta']c_{\beta}a_{\alpha\beta}.$$

Let $n = 0$ in (2.2), we have

$$(2.7) \quad q[\beta'] + a_{\alpha'\beta'} = q[\beta] + a_{\alpha\beta}.$$

• If $a_{\alpha\beta} = 0$, then $[\beta] = q^{2\alpha+1}$ by (2.1). Hence,

$$q^{2\alpha+1} + (q^4 - 1)[\beta] = q^{2\alpha+5} \neq 0.$$

• If $a_{\alpha\beta} \neq 0$ and $[\beta'] \neq 0$, then, by (2.1) and (2.6), we get

$$(2.8) \quad q^{2\alpha'} - q^{-1}[\beta'] = q[\beta].$$

It follows from (2.7) and (2.8) that

$$(2.9) \quad q^{2\alpha} - q^{-1}[\beta] = q[\beta'].$$

Using (2.8) and (2.9), we have that

$$\begin{aligned} 0 \neq q^{2\alpha'} &= q^{-1}[\beta'] + q[\beta] \\ &= q^{-2}(q^{2\alpha} - q^{-1}[\beta]) + q[\beta] \\ &= q^{-3}(q^{2\alpha+1} + (q^4 - 1)[\beta]), \end{aligned}$$

so $q^{2\alpha+1} + (q^4 - 1)[\beta] \neq 0$.

• If $a_{\alpha\beta} \neq 0$ and $[\beta'] = 0$, then $b_{\alpha'\beta'} \neq 0$ by (2.4). It follows from (2.5) that $c_{\beta} = -1$, i.e. $[\beta] = 0$. Hence,

$$q^{2\alpha+1} + (q^4 - 1)[\beta] = q^{2\alpha+1} \neq 0. \quad \blacksquare$$

PROPOSITION 2.6. Let φ be a $\mathbb{Z}$ -grading preserving linear map and $\alpha, \beta, \alpha', \beta' \in \mathbb{C}$. Then $\varphi : A(\alpha, \beta) \simeq A(\alpha', \beta')$ if and only if one of the following conditions holds:

- (1) $q^{2\alpha} = q^{2\alpha'}$ and $[\beta] = [\beta']$;
- (2) $(q^{2\alpha} - 1)(q^{2\alpha'} - 1) \neq 0$ and $[\beta] = [\beta'] = 0$;
- (3) $q^{2\alpha+1} = [\beta] = q^{2\alpha'-1}$, $[\beta'] = 0$ and $q^{2\alpha'} \notin \{q^{2k} \mid k \in \mathbb{Z}\}$;
- (4) $q^{2\alpha'+1} = [\beta'] = q^{2\alpha-1}$, $[\beta] = 0$ and $q^{2\alpha} \notin \{q^{2k} \mid k \in \mathbb{Z}\}$.

PROOF: Assume that $\varphi : A(\alpha, \beta) \simeq A(\alpha', \beta')$. Let

$$\begin{aligned} A(\alpha, \beta) &:= \bigoplus_{k \in \mathbb{Z}} \mathbb{C} v_k, & A(\alpha', \beta') &:= \bigoplus_{k \in \mathbb{Z}} \mathbb{C} v'_k, \\ \varphi(v_k) &:= a_k v'_k, & \text{where } k \in \mathbb{Z} \text{ and } a_k \in \mathbb{C}^*, \end{aligned}$$

and $a_{\alpha\beta}, b_{\alpha\beta}$ defined by (2.1).

Using (1.15), $\varphi d_n(v_k) = d_n \varphi(v_k)$ and the same argument as the one in the proof of Proposition 2.5, we have

$$(2.10) \quad (b_{\alpha\beta}x + q[\beta])a_k = (b_{\alpha'\beta'}x + q[\beta'])a_0,$$

$$(2.11) \quad \begin{aligned} &(-x + q[\beta]y + a_{\alpha\beta})(b_{\alpha\beta}x + q[\beta])(b_{\alpha'\beta'}x + q[\beta']y) \\ &= (-x + q[\beta']y + a_{\alpha'\beta'})(b_{\alpha'\beta'}x + q[\beta'])(b_{\alpha\beta}x + q[\beta]y), \end{aligned}$$

where $x := q^{-2k}$ and $y := q^{2n}$.

Comparing the coefficients of xy , y^2 , x and y in (2.11), we get

$$(2.12) \quad q[\beta]^2 b_{\alpha'\beta'} + [\beta'] a_{\alpha\beta} b_{\alpha\beta} = q[\beta']^2 b_{\alpha\beta} + [\beta] a_{\alpha'\beta'} b_{\alpha'\beta'},$$

$$(2.13) \quad [\beta]^2 [\beta'] = [\beta']^2 [\beta],$$

$$(2.14) \quad [\beta] a_{\alpha\beta} b_{\alpha'\beta'} = [\beta'] a_{\alpha'\beta'} b_{\alpha\beta},$$

$$(2.15) \quad [\beta][\beta'] a_{\alpha\beta} = [\beta][\beta'] a_{\alpha'\beta'}.$$

- If $[\beta][\beta'] \neq 0$, then (1) follows from (2.13) and (2.15).
- If $[\beta] = [\beta'] = 0$, then, by (2.10)

$$(q^{2\alpha} - 1)a_k = (q^{2\alpha'} - 1)a_0.$$

So $q^{2\alpha} - 1 = 0 \Leftrightarrow q^{2\alpha'} - 1 = 0$, i.e., either (1) holds, or (2) holds.

- If $[\beta] \neq 0$ and $[\beta'] = 0$, then $[\beta] a_{\alpha\beta} b_{\alpha'\beta'} = 0$ by (2.14). Since $b_{\alpha'\beta'} \neq 0$ by (2.10), $a_{\alpha\beta} = 0$. Hence,

$$(2.16) \quad q^{2\alpha+1} = [\beta] = q^{2\alpha'-1},$$

where the last equation in (2.16) follows from (2.12).

By (2.10), we have

$$(-q^{-2k} + q^{2\alpha'})a_k = (b_{\alpha\beta}x + q[\beta])a_k = b_{\alpha'\beta'}x a_0 \neq 0$$

for all $k \in \mathbb{Z}$. This implies that

$$(2.17) \quad q^{2\alpha'} \notin \{q^{2k} \mid k \in \mathbb{Z}\}.$$

(2.16) and (2.17) implies that (3) in Proposition 2.6 holds.

• If $[\beta] = 0$ and $[\beta'] \neq 0$, then $[\beta']a_{\alpha'\beta'}b_{\alpha\beta} = 0$ by (2.14). Since $b_{\alpha\beta} \neq 0$ by (2.10), $a_{\alpha'\beta'} = 0$. Hence,

$$(2.18) \quad q^{2\alpha'+1} = [\beta'] = q^{2\alpha-1},$$

where the last equation in (2.18) follows from (2.12).

It follows from (2.10) that

$$q^{2\alpha} \notin \{q^{2k} \mid k \in \mathbb{Z}\}.$$

By (2.18) and (2.19), (4) in Proposition 2.6 holds.

Conversely, if one of the conditions in Proposition 2.6 holds, then the $\mathbb{Z}$ -grading preserving linear map φ defined by

$$\varphi(v_k) := \frac{(q^{2\alpha'} - q^{-1}[\beta'] - 1)q^{-2k} + q[\beta']}{(q^{2\alpha} - q^{-1}[\beta] - 1)q^{-2k} + q[\beta]} v'_k \quad \text{for } k \in \mathbb{Z}$$

is an isomorphism from $A(\alpha, \beta)$ to $A'(\alpha', \beta')$. ■

PROPOSITION 2.7. Let φ be a $\mathbb{Z}$ -grading preserving linear map and $\alpha, \beta, \alpha', \beta' \in \mathbb{C}$. Then $\varphi : B(\alpha, \beta) \simeq B(\alpha', \beta')$ if and only if one of the following conditions holds:

- (1) $q^{2\alpha} = q^{2\alpha'}$ and $[\beta] = [\beta']$;
- (2) $q^{2\alpha} = q^{-1}[\beta]$, $q^{2\alpha'} = q^{-1}[\beta']$ and $(q[\beta] - 1)(q[\beta'] - 1) \neq 0$;
- (3) $q^{2\alpha+1} = [\beta] = q^{2\alpha'-1}$, $[\beta'] = 0$ and $q^{2\alpha'} \notin \{q^{2k} \mid k \in \mathbb{Z}\}$;
- (4) $q^{2\alpha'+1} = [\beta] = q^{2\alpha-1}$, $[\beta] = 0$ and $q^{2\alpha} \notin \{q^{2k} \mid k \in \mathbb{Z}\}$.

PROOF: Assume that $\varphi : B(\alpha, \beta) \simeq B(\alpha', \beta')$. Let

$$\begin{aligned} B(\alpha, \beta) &:= \bigoplus_{k \in \mathbb{Z}} \mathbb{C} v_k, & B(\alpha', \beta') &:= \bigoplus_{k \in \mathbb{Z}} \mathbb{C} v'_k, \\ \varphi(v_k) &:= a_k v'_k, & \text{where } k \in \mathbb{Z} \text{ and } a_k \in \mathbb{C}^*, \end{aligned}$$

and $a_{\alpha\beta}, c_\beta$ are defined by (2.1).

Using (1.16), $\varphi d_n(v_k) = d_n \varphi(v_k)$ and the same argument as the one in the proof of Proposition 2.5, we have

$$(2.20) \quad (c_\beta x + a_{\alpha\beta}) a_k = (c_{\beta'} x + a_{\alpha'\beta'}) a_0.$$

$$(2.21) \quad \begin{aligned} & (-x + a_{\alpha\beta}y + q[\beta])(c_{\beta}x + a_{\alpha\beta})(c_{\beta'}x + a_{\alpha'\beta'}y) \\ & = (-x + a_{\alpha'\beta'}y + q[\beta'])(c_{\beta'}x + a_{\alpha'\beta'})(c_{\beta}x + a_{\alpha\beta}y). \end{aligned}$$

Comparing the coefficients of x^2 , xy , y^2 , x and y in (2.21), we get

$$(2.22) \quad -c_{\beta'}a_{\alpha\beta} + q[\beta]c_{\beta}c_{\beta'} = -c_{\beta}a_{\alpha'\beta'} + q[\beta']c_{\beta}c_{\beta'},$$

$$(2.23) \quad c_{\beta'}a_{\alpha\beta}^2 + q[\beta]c_{\beta}a_{\alpha'\beta'} = c_{\beta}a_{\alpha'\beta'}^2 + q[\beta']c_{\beta'}a_{\alpha\beta},$$

$$(2.24) \quad a_{\alpha\beta}^2a_{\alpha'\beta'} = a_{\alpha'\beta'}^2a_{\alpha\beta},$$

$$(2.25) \quad [\beta]a_{\alpha\beta}c_{\beta'} = [\beta']a_{\alpha'\beta'}c_{\beta},$$

$$(2.26) \quad [\beta]a_{\alpha\beta}a_{\alpha'\beta'} = [\beta']a_{\alpha\beta}a_{\alpha'\beta'}.$$

- If $a_{\alpha\beta}a_{\alpha'\beta'} \neq 0$, then (1) in Proposition 2.7 follows from (2.24) and (2.26).
- If $a_{\alpha\beta} = a_{\alpha'\beta'} = 0$, then, by (2.20)

$$(q[\beta] - 1)a_k = (q[\beta'] - 1)a_0.$$

So $(q[\beta] - 1) = 0 \Leftrightarrow q[\beta'] - 1 = 0$, i.e., either (1) holds, or (2) holds.

- If $a_{\alpha\beta} = 0$ and $a_{\alpha'\beta'} \neq 0$, then

$$(2.20) \Rightarrow c_{\beta} \neq 0.$$

$$(2.25) \Rightarrow [\beta'] = 0.$$

$$(2.23) \Rightarrow q[\beta]c_{\beta}a_{\alpha'\beta'} = c_{\beta}a_{\alpha'\beta'}^2 \Rightarrow q[\beta] = a_{\alpha'\beta'} = q^{2\alpha'} \Rightarrow [\beta] = q^{2\alpha'-1}.$$

$$a_{\alpha\beta} = 0 \Rightarrow [\beta] = q^{2\alpha+1}.$$

Furthermore, by (2.20)

$$0 \neq c_{\beta}xa_k = (c_{\beta'}x + a_{\alpha'\beta'})a_0 = (-q^{-2k} + q^{2\alpha'})a_0 \Rightarrow q^{2\alpha'} \neq q^{-2k},$$

where $k \in \mathbb{Z}$, i.e.,

$$q^{2\alpha'} \notin \{q^{2k} \mid k \in \mathbb{Z}\}.$$

So (3) holds in this case.

- If $a_{\alpha\beta} \neq 0$ and $a_{\alpha'\beta'} = 0$, then

$$(2.20) \Rightarrow c_{\beta'} \neq 0.$$

$$(2.25) \Rightarrow [\beta'] = 0.$$

$$(2.23) \Rightarrow c_{\beta'}a_{\alpha\beta}^2 = q[\beta']c_{\beta'}a_{\alpha\beta} \Rightarrow q^{2\alpha} = a_{\alpha\beta} = q[\beta'] \Rightarrow [\beta'] = q^{2\alpha-1}.$$

$$a_{\alpha'\beta'} = 0 \Rightarrow q^{2\alpha} \notin \{q^{2k} \mid k \in \mathbb{Z}\}.$$

$$(2.20) \Rightarrow q^{2\alpha} \notin \{q^{2k} \mid k \in \mathbb{Z}\}.$$

Hence, (4) is true.

Conversely, if one of the conditions in Proposition 2.7 holds, then the $\mathbb{Z}$ -grading preserving linear map φ defined by

$$\varphi(v_k) := \frac{(q[\beta'] - 1)q^{-2k} + (q^{2\alpha'} - q^{-1}[\beta'])}{(q[\beta] - 1)q^{-2k} + (q^{2\alpha} - q^{-1}[\beta])} v'_k \quad \text{for } k \in \mathbb{Z}$$

is an isomorphism from $B(\alpha, \beta)$ to $B(\alpha', \beta')$. ■

PROPOSITION 2.7. For $(\alpha, \beta) \in \mathbb{C} \times \mathbb{C}$, we have

- (1) $A(\alpha, \beta)$ is reducible $\Leftrightarrow$ either $q^{2\alpha} - q^{2t} = [\beta] = 0$ or $q^{2\alpha+1} = [\beta] = q^{-2t-1}$ for some $t \in \mathbb{Z}$.
- (2) If $B(\alpha, \beta) \notin \text{cl}(A)$, then $B(\alpha, \beta)$ is irreducible.

PROOF: (1). $\Rightarrow$: Let N be a non-zero submodule of $A(\alpha, \beta)$ with $N \neq A(\alpha, \beta)$. Then N is a direct sum of some of the 1-dimensional subspace $\mathbb{C}v_j$. Let $v_t \in N$.

• If there exist two distinct vectors v_m and v_n ($m \neq n$) in $A(\alpha, \beta) \setminus N$, then, by (1.15)

$$\begin{aligned} N \ni d_{m-t}(v_t) &= -\frac{q^t}{q - q^{-1}}(-q^{-2t} + q[\beta]q^{2(m-t)} + (q^{2\alpha} - q^{-1}[\beta]))v_m, \\ N \ni d_{n-t}(v_t) &= -\frac{q^t}{q - q^{-1}}(-q^{-2t} + q[\beta]q^{2(n-t)} + (q^{2\alpha} - q^{-1}[\beta]))v_n. \end{aligned}$$

Hence,

$$(2.27) \quad -q^{-2t} + q[\beta]q^{2(m-t)} + (q^{2\alpha} - q^{-1}[\beta]) = 0,$$

$$(2.28) \quad -q^{-2t} + q[\beta]q^{2(n-t)} + (q^{2\alpha} - q^{-1}[\beta]) = 0.$$

It follows from (2.27)–(2.28) that $[\beta] = 0$. Going back to (2.27), we get that $q^{2\alpha} = q^{-2t}$ and t is unique. In other words, $N = \mathbb{C}v_t$ in this case.

• If $N = \bigoplus_{\substack{k \in \mathbb{Z} \\ k \neq t}} \mathbb{C}v_k$, then we choose $v_m, v_n \in N$ with $m \neq n$. By (1.15),

$$\begin{aligned} N \ni d_{t-m}(v_m) &= -\frac{q^m}{q - q^{-1}}(-q^{-2m} + q[\beta]q^{2(t-m)} + (q^{2\alpha} - q^{-1}[\beta]))v_t, \\ N \ni d_{t-n}(v_n) &= -\frac{q^n}{q - q^{-1}}(-q^{-2n} + q[\beta]q^{2(t-n)} + (q^{2\alpha} - q^{-1}[\beta]))v_t. \end{aligned}$$

Since $v_t \notin N$, we have to have

$$(2.29) \quad -q^{-2m} + q[\beta]q^{2(t-m)} + (q^{2\alpha} - q^{-1}[\beta]) = 0,$$

$$(2.30) \quad -q^{-2n} + q[\beta]q^{2(t-n)} + (q^{2\alpha} - q^{-1}[\beta]) = 0.$$

It follows from (2.29)–(2.30) that

$$(q^{-2m} - q^{-2n})(q^{2t+1}[\beta] - 1) = 0,$$

or

$$q^{2t+1}[\beta] = 1.$$

Going back to (2.29), we get

$$q^{2\alpha+1} = [\beta].$$

(1). $\Leftarrow$: If $q^{2\alpha} - q^{-2t} = [\beta] = 0$, then $\mathbb{C}v_t$ is a submodule of $A(\alpha, \beta)$.
If $q^{2\alpha+1} = [\beta] = q^{-2t-1}$, then $\bigoplus_{\substack{k \in \mathbb{Z} \\ k \neq t}} \mathbb{C}v_k$ is a submodule of $A(\alpha, \beta)$.

Therefore, $A(\alpha, \beta)$ is always reducible in these cases.

(2). If $B(\alpha, \beta) \notin \text{cl}(A)$, then, by Proposition 2.5,

$$(2.31) \quad q^{2\alpha+1} + (q^4 - 1)[\beta] = 0.$$

Assume that $N \neq 0$ is a submodule of $B(\alpha, \beta)$, it follows from Proposition 2.4 that $N = \bigoplus_{k \in S} \mathbb{C}v_k$ for some non-empty subset S of $\mathbb{Z}$.

• If there exists $v_m \notin N$ and $v_n \notin N$ with $m \neq n$, then by (1.16),

$$N \ni d_{m-k}(v_k) = -\frac{q^k}{q - q^{-1}} \left(-q^{-2k} + (q^{2\alpha} - q^{-1}[\beta])q^{2(m-k)} + q[\beta] \right) v_m,$$

$$N \ni d_{n-k}(v_k) = -\frac{q^k}{q - q^{-1}} \left(-q^{-2k} + (q^{2\alpha} - q^{-1}[\beta])q^{2(n-k)} + q[\beta] \right) v_n,$$

where $v_k \in N$. So the coefficients of v_m and v_n have to be zero. This implies that $q^{2\alpha} - q^{-1}[\beta] = 0$. Going back to (2.31), we get that $[\beta] = 0$, which contradicts (2.31).

• If $\mathbb{Z} \setminus S = \{s\}$, then we can choose $v_m \in N$ and $v_n \in N$ with $m \neq n$. As above, it follows from $d_{s-m}(v_m) \in N$ and $d_{s-n}(v_n) \in N$ that $[\beta] = 0$, which is impossible. ■

For $t \in \mathbb{Z}$, we define

$$A_t(\alpha, \beta) = \begin{cases} \frac{A(\alpha, \beta)}{\mathbb{C}v_t}, & \text{if } q^{2\alpha} - q^{-2t} = [\beta] = 0; \\ \bigoplus_{\substack{k \in \mathbb{Z} \\ k \neq t}} \mathbb{C}v_k, & \text{if } q^{2\alpha+1} = [\beta] = q^{-2t-1}. \end{cases}$$

Then, $A_t(\alpha, \beta)$ is an irreducible $U(W_q)$ -module by the proof of (1) in Proposition 2.7.

PROPOSITION 2.8. Let $q \in \mathbb{R}$, then with respect to the antilinear anti-involution θ of $U(W_q)$,

- (1) $A(\alpha, \beta)$ is unitary $\iff q > 0$ and $q^{2\alpha} = q^{-1}[\beta] + q[\bar{\beta}]$.
- (2) $A_t(\alpha, \beta)$ and $B(\alpha, \beta)$ are not unitary, where $B(\alpha, \beta) \notin \text{cl}(A)$.

PROOF: (1). $\Rightarrow$: Assume that $A(\alpha, \beta) = \oplus_{k \in \mathbb{Z}} \mathbb{C} v_k$ is unitary and $\langle \cdot, \cdot \rangle$ the contravariant form on V . So

$$\langle d_m(v_k) | v_\ell \rangle = \langle v_k | d_{-m}(v_\ell) \rangle \quad \text{for } m, k, \ell \in \mathbb{Z}.$$

Let $n := \ell := m + k$, then we have, by (1.15)

$$\begin{aligned} & q^k (-q^{-2k} + q[\beta]q^{2n-2k} + (q^{2\alpha} - q^{-1}[\beta])) \langle v_n | v_n \rangle \\ (2.32) \quad & = q^n (-q^{-2n} + q[\bar{\beta}]q^{-2n+2k} + (q^{2\alpha} - q^{-1}[\bar{\beta}])) \langle v_k | v_k \rangle, \end{aligned}$$

where $n, k \in \mathbb{Z}$. Let $k = 0$ in (2.32), we get

$$(2.33) \quad (q[\beta]q^{2n} + b_{\alpha\beta}) \langle v_n | v_n \rangle = q^n (\bar{c}_\beta q^{-2n} + \bar{a}_{\alpha\beta}) \langle v_0 | v_0 \rangle,$$

where $a_{\alpha\beta}$, $b_{\alpha\beta}$ and c_β are defined by (2.1).

It follows from (2.32) and (2.33) that

$$\begin{aligned} & (-x + q[\beta]xy + a_{\alpha\beta})(b_{\alpha\beta}x + q[\beta])(\bar{a}_{\alpha\beta}y + \bar{c}_\beta) \\ (2.34) \quad & = (-x + \bar{a}_{\alpha\beta}xy + q[\bar{\beta}])(\bar{c}_\beta x + \bar{a}_{\alpha\beta})(q[\beta]y + b_{\alpha\beta}), \end{aligned}$$

where $x := q^{-2k}$ and $y := q^{2n}$. Comparing the coefficients of x^2y , xy and y , we get

$$(2.35) \quad -\bar{a}_{\alpha\beta}b_{\alpha\beta} + q[\beta]\bar{c}_\beta b_{\alpha\beta} = -q[\beta]\bar{c}_\beta + \bar{c}_\beta \bar{a}_{\alpha\beta} b_{\alpha\beta},$$

$$(2.36) \quad q^2[\beta]^2 \bar{c}_\beta + a_{\alpha\beta} \bar{a}_{\alpha\beta} b_{\alpha\beta} = \bar{a}_{\alpha\beta}^2 b_{\alpha\beta} + q^2[\beta][\bar{\beta}] \bar{c}_\beta,$$

$$(2.37) \quad [\beta]a_{\alpha\beta} \bar{a}_{\alpha\beta} = q[\beta][\bar{\beta}] \bar{a}_{\alpha\beta}.$$

Suppose that $[\beta] = 0$, then $a_{\alpha\beta} = q^{2\alpha} \neq 0$, $c_\beta = -1$ and $b_{\alpha\beta} \neq 0$ by (2.1) and (2.33). It follows from (2.36) that $a_{\alpha\beta} = \bar{a}_{\alpha\beta}$. So (2.33) becomes that

$$(q^{2\alpha} - 1) \langle v_n | v_n \rangle = q^n (q^{2\alpha} - q^{-2n}) \langle v_0 | v_0 \rangle \quad \text{for } n \in \mathbb{Z}.$$

This implies that $f(n) := \frac{q^{2n} - q^{4n}}{q^{2n} - 1} > 0$ for all $n \in \mathbb{Z}$, which is impossible because $f(n)f(-n) < 0$ for large $n > 0$. Therefore, we have proved that $[\beta] \neq 0$.

Similarly, we can prove that $a_{\alpha\beta} \neq 0$ by using (2.35).

Going back to (2.37), we have $a_{\alpha\beta} = q[\overline{\beta}]$, i.e. $q^{2\alpha} = q^{-1}[\beta] + q[\overline{\beta}]$.

Finally, choose an odd $n_0 \in \mathbb{Z}$ such that $q[\beta]q^{2n_0} + b_{\alpha\beta} \neq 0$, then (2.33) gives that

$$\langle v_{n_0} | v_{n_0} \rangle = q^{-n_0} \langle v_0 | v_0 \rangle,$$

which implies that $q > 0$.

⇐: Define a Hermitian form $\langle \cdot | \cdot \rangle$ on V by

$$\langle v_n | v_m \rangle := \delta_{nm} q^{-n} \quad \text{for all } n, m \in \mathbb{Z}.$$

It is easy to check that $\langle \cdot | \cdot \rangle$ is a contravariant form.

(2). Assume that $A_t(\alpha, \beta)$ is unitary, then, using the same argument as above, we can get

$$q^{2\alpha} = q^{-1}[\beta] + q[\overline{\beta}],$$

which is impossible because we have either $q^{2\alpha} - q^{2t} = [\beta] = 0$ or $q^{2\alpha+1} = [\beta] = q^{-2t-1}$. This proves that $A_t(\alpha, \beta)$ is not unitary.

Finally, we prove that if $B(\alpha, \beta) \notin \text{cl}(A)$, then $B(\alpha, \beta)$ is not unitary.

Suppose that $B(\alpha, \beta)$ has a contravariant form $\langle \cdot | \cdot \rangle$ and $\langle v | v \rangle > 0$ for $0 \neq v \in B$, then

$$(2.38) \quad \langle d_m(v_k) | v_\ell \rangle = \langle v_k | d_{-m}(v_\ell) \rangle \quad \text{for } m, k, \ell \in \mathbb{Z}.$$

By (1.16) and Proposition 2.5,

$$(2.39) \quad d_n(v_j) = -\frac{q^k}{q - q^{-1}}(-q^{-2k} + q^3[\beta]q^{2n} + q[\beta])v_{n+j} \quad \text{for } n, j \in \mathbb{Z}.$$

Let $n := \ell := m + k$, then, by (2.38) and (2.39)

$$\begin{aligned} & q^k(-q^{-2k} + q^3[\beta]q^{2n-2k} + q[\beta]) \langle v_n | v_n \rangle \\ &= q^n(-q^{-2n} + q^3[\overline{\beta}]q^{-2n+2k} + q[\overline{\beta}]) \langle v_k | v_k \rangle. \end{aligned}$$

Let $n = k$ in (2.40), we get

$$q^3[\beta] + q[\beta] = q^3[\overline{\beta}] + q[\overline{\beta}] \quad \text{or} \quad [\beta] = [\overline{\beta}].$$

Hence, (2.40) becomes

$$(2.41) \quad \begin{aligned} q^k(-q^{-2k} + q^3[\beta]q^{2n-2k} + q[\beta]) &\leq v_n |v_n| > \\ &= q^n(-q^{-2n} + q^3[\beta]q^{-2n+2k} + q[\beta]) \leq v_k |v_k| > . \end{aligned}$$

Using (2.41) and the same argument as the one in the proof of (1), we have

$$\begin{aligned} &(-x + q^3[\beta]xy + q[\beta])(q[\beta]y + (q^3[\beta] - 1)((q[\beta] - 1)x + q^3[\beta])) \\ &= (q[\beta]xy - x + q^3[\beta])((q^3[\beta] - 1)x + q[\beta])(q^3[\beta]y + (q[\beta] - 1)), \end{aligned}$$

where $x := q^{-2k}$ and $y := q^{2n}$. Comparing the constants in the equation above, we conclude that

$$q[\beta] \cdot (q^3[\beta] - 1) \cdot q^3[\beta] = q^3[\beta] \cdot q[\beta] \cdot (q[\beta] - 1).$$

Since $[\beta] \neq 0$, $q^3 = q$, i.e., $q^2 = 1$, which is impossible. ■

§3.3. q -analogue of Kaplansky's theorem

LEMMA 3.1. *Let q be not a root of unity, then for all integers n and all positive integers s , we have in $U(W_q)$*

$$\begin{aligned} d_n d_{-n}^s &= q^{-2ns} d_{-n}^s d_n + q^{-sn} \frac{[2n][sn]}{[n]} d_{-n}^{s-1} d_0 J^{-1} \\ &\quad + [sn][(s-1)n] d_{-n}^{s-1} J^{-2}. \end{aligned}$$

PROOF: We use induction on s . It is clear that the Lemma is true for $s = 1$. Now we assume that the Lemma is true for s , then

$$\begin{aligned} d_n d_{-n}^{s+1} &= (d_n d_{-n}^s) d_{-n} \\ &= (q^{-2ns} d_{-n}^s d_n + q^{-sn} \frac{[2n][sn]}{[n]} d_{-n}^{s-1} d_0 J^{-1} \\ &\quad + [sn][(s-1)n] d_{-n}^{s-1} J^{-2}) d_{-n} \\ &= q^{-2ns} d_{-n}^s (q^{-2n} d_{-n} d_n + q^{-n} [2n] d_0 J^{-1}) \\ &\quad + [sn][(s-1)n] q^{2n} d_{-n}^s J^{-2} \\ &\quad + q^{-sn} \frac{[2n][sn]}{[n]} d_{-n}^{s-1} q^n (q^{-n} d_{-n} d_0 + [n] d_{-n} J^{-1}) J^{-1} \\ &= q^{-2n(s+1)} d_{-n}^{s+1} d_n + q^{-(s+1)n} (q^{-sn} [n] + q[sn]) \frac{[2n]}{[n]} d_{-n}^s d_0 J^{-1} \\ &\quad + (q^{-(s-1)n} [2n] + q^{2n} [(s-1)] [sn] d_{-n}^s J^{-2} \\ &= q^{-2n(s+1)} d_{-n}^{s+1} d_n + q^{-(s+1)n} \frac{[(s+1)n][2n]}{[n]} d_{-n}^s d_0 J^{-1} \\ &\quad + [(s+1)n][sn] d_{-n}^s J^{-2}. \end{aligned}$$

This proves the Lemma. ■

THEOREM 3.2. Let q be not a root of unity and $V = \bigoplus_{k \in \mathbb{Z}} \mathbb{C}v_k$ a $\mathbb{Z}$ -graded $U(W_q)$ -module with $J(v_k) \in \mathbb{C}v_k$ for $k \in \mathbb{Z}$. If d_1 and d_{-1} are injective operators on V and

$$(Jd_1d_{-1}J - Jd_{-1}d_1J)(v_0) \neq \frac{1}{q - q^{-1}}v_0,$$

then $V \simeq A(\lambda, \alpha, \beta)$ or $V \simeq B(\lambda, \alpha, \beta)$ for some $(\lambda, \alpha, \beta) \in \mathbb{C}^* \times \mathbb{C} \times \mathbb{C}$.

PROOF: Since $J(v_k) \in \mathbb{C}v_k$, $d_1(v_k) \neq 0$ and $Jd_1J^{-1}(v_k) = qd_1(v_k)$, there exists some $\lambda \in \mathbb{C}^*$ such that $J(v_k) = \lambda q^k v_k$ for all $k \in \mathbb{Z}$. Using the automorphism $\varphi(\lambda^{-1})$, we can assume that $\lambda = 1$, in which case, we will prove that either $V \simeq A(\alpha, \beta)$ or $V \simeq B(\alpha, \beta)$ for some $(\alpha, \beta) \in \mathbb{C} \times \mathbb{C}$.

Set

$$d_0(v_0) = av_0, \quad d_1d_{-1}(v_j) = x_jv_j, \quad d_{-1}d_1(v_j) = y_jv_j,$$

where $a, x_j, y_j \in \mathbb{C}$ and $j \in \mathbb{Z}$. We consider the system (i) with respect to α and β :

$$(3.1) \quad -([\alpha]q^\alpha + [\beta]) = a,$$

$$(3.2) \quad [\alpha]q^\alpha([\alpha - 1]q^\alpha + [2][\beta]) = x_0$$

and the system (ii) with respect to α and β :

$$(3.3) \quad \begin{aligned} & -([\alpha]q^\alpha + [\beta]) = a, \\ & [\alpha + 1]q^\alpha([\alpha]q^\alpha + [2][\beta]q) = y_0. \end{aligned}$$

First, we assume that there exist α and β such that (i) holds. Using induction on j and (3.1) gives us

$$(3.4) \quad d_0(v_j) = -([\alpha + j]q^\alpha + [\beta]q^j)v_j \quad \text{for } j \in \mathbb{Z}.$$

Since $(qd_1d_{-1}J - q^{-1}d_{-1}d_1J)(v_j) = [2]d_0v_j$, we have by (3.4)

$$(3.5) \quad q^{j+1}x_j - q^{j-1}y_j = -[2][\alpha + j]q^\alpha - [2][\beta]q^j \quad \text{for } j \in \mathbb{Z}.$$

Furthermore, computing $d_1d_{-1}d_1(v_{j-1})$ in two ways produces the following relation between x_j and y_j :

$$(3.6) \quad x_j = y_{j-1} \quad \text{for } j \in \mathbb{Z}.$$

Going back to (3.5), we get

$$(3.7) \quad q^{j+1}x_j - q^{j-1}x_{j+1} = -[2][\alpha + j]q^\alpha - [2][\beta]q^j \quad \text{for } j \in \mathbb{Z}.$$

Now we claim that

$$(3.8) \quad x_j = [\alpha + j]q^\alpha([\alpha + j - 1]q^\alpha + [2][\beta]q^j) \quad \text{for } j \in \mathbb{Z}.$$

By (3.2), (3.8) is true for $j = 0$. Assume that (3.8) is true for j , then (3.8) is also true for $j \pm 1$. For example, let us prove that (3.8) is true for $j + 1$. By (3.7),

$$\begin{aligned} x_{j+1} &= q^2 x_j + [2][\alpha + j]q^{\alpha-j+1} + [2][\beta]q \\ &= q^2[\alpha + j]q^\alpha([\alpha + j - 1]q^\alpha + [2][\beta]q^j) + [2][\alpha + j]q^{\alpha-j+1} + [2][\beta]q \\ &= [\alpha + j]q^{2\alpha}(q^2[\alpha + j - 1] + [2]q^{-\alpha-j+1}) \\ &\quad + [2][\beta]q^\alpha([\alpha + j]q^{2+j} + q^{1-\alpha}) \\ &= [\alpha + j]q^{2\alpha}[\alpha + j + 1] + [2][\beta]q^\alpha q^{j+1}[\alpha + j + 1] \\ &= [\alpha + j + 1]q^\alpha([\alpha + j]q^\alpha + [2][\beta]q^{j+1}). \end{aligned}$$

Hence, (3.8) is true for all $j \in \mathbb{Z}$ by induction.

Let $j = 1$ in (3.8), we get (3.3). So we have proved that if α and β satisfy (i), then α and β also satisfy (ii).

Similarly, we can prove that if α and β satisfy (ii), then α and β also satisfy (i).

Now we prove that either (i) has a solution or (ii) has a solution.

Choose $\beta, \beta' \in \mathbb{C}$ such that

$$(3.9) \quad (-a - [\beta])((-a - [\beta])q^{-1} - 1 + [2][\beta]) = x_0,$$

$$(3.10) \quad ((-a - [\beta'])q + 1)((-a - [\beta']) + [2][\beta']q) = y_0.$$

Suppose that both (i) and (ii) has no solution, then we have to have

$$(3.11) \quad -\left(\frac{q^{2\alpha} - 1}{q - q^{-1}} + [\beta]\right) \neq a \quad \text{for all } \alpha \in \mathbb{C};$$

$$(3.12) \quad -\left(\frac{q^{2\alpha'} - 1}{q - q^{-1}} + [\beta']\right) \neq a \quad \text{for all } \alpha' \in \mathbb{C}.$$

In fact, if $-\left(\frac{q^{2\alpha} - 1}{q - q^{-1}} + [\beta]\right) = a$ for some $\alpha \in \mathbb{C}$, then α and β satisfies (3.1). Hence, by (3.9)

$$\begin{aligned} x_0 &= [\alpha]q^\alpha([\alpha]q^\alpha \cdot q^{-1} - 1 + [2][\beta]) \\ &= [\alpha]q^\alpha([\alpha - 1]q^\alpha + [2][\beta]), \end{aligned}$$

i.e., α and β also satisfies (3.2). So (α, β) is a solution of (i), which contradicts our assumption. This proves (3.11).

Similarly, if $-\left(\frac{q^{2\alpha'} - 1}{q - q^{-1}} + [\beta']\right) = a$ for some $\alpha' \in \mathbb{C}$, then

$$-([\alpha']q^{\alpha'} + [\beta']) = a$$

and, by (3.10)

$$\begin{aligned} y_0 &= ([\alpha']q^{\alpha'} \cdot q + 1)([\alpha']q^{\alpha'} + [2][\beta']q) \\ &= [\alpha' + 1]q^{\alpha'}([\alpha']q^{\alpha'} + [2][\beta']q). \end{aligned}$$

Hence, (α', β') is a solution of (ii), which is impossible by our assumption. So (3.12) holds.

It follows from (3.11) and (3.12) that

$$(3.13) \quad [\beta] = [\beta'] = \frac{1}{q - q^{-1}} - a.$$

By (3.9) and (3.13), we have

$$\begin{aligned} x_0 &= (-a - [\beta])(-aq^{-1} - 1 + ([2] - q^{-1})[\beta]) \\ &= (-a - [\beta])(-aq^{-1} - 1 + q[\beta]) \\ &= -\frac{1}{q - q^{-1}} \left(-aq^{-1} - 1 + q \left(\frac{1}{q - q^{-1}} - a \right) \right) \\ &= -\frac{1}{q - q^{-1}} \left(-a(q + q^{-1}) + \left(-1 + \frac{q}{q - q^{-1}} \right) \right) \\ &= -\frac{1}{q - q^{-1}} \left(-a(q + q^{-1}) + \frac{q^{-1}}{q - q^{-1}} \right) \\ &= \frac{1}{(q - q^{-1})^2} (a(q^2 - q^{-2}) - q^{-1}), \end{aligned}$$

i.e.,

$$(3.14) \quad a(q^2 - q^{-2}) - q^{-1} = (q - q^{-1})^2 x_0.$$

By (3.10) and (3.13), we have

$$\begin{aligned} y_0 &= ((-a - [\beta'])q + 1)(-a + ([2]q - 1)[\beta']) \\ &= \left(-\frac{q}{q - q^{-1}} + 1 \right) \left(-a + q^2 \left(\frac{1}{q - q^{-1}} - a \right) \right) \\ &= \frac{q^{-1}}{q - q^{-1}} \left(a(q^2 + 1) - \frac{q^2}{q - q^{-1}} \right) \\ &= \frac{1}{(q - q^{-1})^2} (a(q^2 - q^{-2}) - q), \end{aligned}$$

i.e.,

$$(3.15) \quad a(q^2 - q^{-2}) - q = (q - q^{-1})^2 y_0.$$

(3.14)–(3.15) gives us

$$q - q^{-1} = (q - q^{-1})^2 (x_0 - y_0),$$

or

$$\frac{1}{q - q^{-1}} = x_0 - y_0,$$

which contradicts the assumption in Theorem 3.2. This proves that either (i) has a solution or (ii) has a solution. Therefore there exists $(\alpha, \beta) \in \mathbb{C} \times \mathbb{C}$ such that (i) holds.

Using (3.4) and (3.8), we can choose a basis of V , say $\{v_k \mid k \in \mathbb{Z}\}$, such that

$$(3.16) \quad d_n(v_j) = -([\alpha + j]q^\alpha + [n + 1][\beta]q^{n+j})v_{n+j},$$

where $n = 0, \pm 1$ and $j \in \mathbb{Z}$.

For $j \in \mathbb{Z}$, set

$$d_2(v_j) := e(j)v_{j+2}, \quad d_{-2}(v_j) := g(j)v_{j-2},$$

$$(3.17) \quad e(j) := f(j) - ([j + \alpha]q^\alpha + [3][\beta]q^{j+2}),$$

$$(3.18) \quad g(j) := h(j) - ([j + \alpha]q^\alpha - [\beta]q^{j-2}),$$

where $e(j), g(j), f(j), h(j) \in \mathbb{C}$.

Using (3.16) and following identities:

$$(q^2 d_2 d_{-1} J - q^{-1} d_{-1} d_2 J)(v_j) = [3]d_1(v_j),$$

$$(q^{-2} d_{-2} d_1 J - q d_1 d_{-2} J)(v_j) = -[3]d_{-1}(v_j),$$

we get

$$q^{\alpha+j-1}[\alpha + j + 2]e(j) - q^{\alpha+j+2}[\alpha + j]e(j-1) = -[3]([\alpha + j]q^\alpha + [2][\beta]q^{j+1}),$$

$$\begin{aligned} & q^{j+1}([\alpha + j - 2]q^\alpha + [2][\beta]q^{j-1})g(j) \\ & - q^{j-2}([\alpha + j]q^\alpha + [2][\beta]q^{j+1})g(j+1) = [3][\alpha + j]q^\alpha. \end{aligned}$$

It follows from (3.17) and (3.18) that

$$\begin{aligned} [\alpha + j + 2]f(j) &= q^3[\alpha + j]f(j-1), \\ ([\alpha + j]q^\alpha + [2][\beta]q^{j+1})h(j+1) &= q^3([\alpha + j - 2]q^\alpha + [2][\beta]q^{j-1})h(j). \end{aligned}$$

These identities imply that

$$(3.19) \quad f(j) = \frac{q^{3j}[\alpha + 1][\alpha + 2]}{[\alpha + j + 1][\alpha + j + 2]}f(0),$$

$$(3.20) \quad h(j) = \frac{q^{3j}([\alpha - 1]q^\alpha + [2][\beta])([\alpha - 2]q^\alpha + [2][\beta]q^{-1})}{([\alpha + j - 2]q^\alpha + [2][\beta]q^{j-1})([\alpha + j - 1]q^\alpha + [2][\beta]q^j)}h(0),$$

where $j \in \mathbb{Z}$. Note, that denominators in (3.19) and (3.20) are non-zero follows from (3.16) and $d_{\pm 1}(v_j) \neq 0$ for all $j \in \mathbb{Z}$.

Let $z := q^{-j}[j]$. We can rewrite (3.17)—(3.20) as follows:

$$(3.21) \quad q^{-j}e(j-2) = q^{-j}f(j-2) - (q^2z + [\alpha - 2]q^\alpha + [3][\beta]).$$

$$(3.22) \quad q^{-j}g(j) = q^{-j}h(j) - (z + [\alpha]q^\alpha - [\beta]q^{-2}).$$

$$(3.23) \quad q^{-j}f(j-2) = \frac{q^{-2}[\alpha + 1][\alpha + 2]f(0)}{(q^{3-\alpha}z + [\alpha + 1] - [2]q^{1-\alpha})(q^{2-\alpha}z + [\alpha + 2] - [2]q^{-\alpha})}.$$

$$(3.24) \quad q^{-j}h(j) = \frac{([\alpha - 1]q^\alpha + [2][\beta])([\alpha - 2]q^\alpha + [2][\beta]q^{-1})h(0)}{(qz + [\alpha - 1]q^\alpha + [2][\beta])(q^2z + [\alpha - 2]q^\alpha + [2][\beta]q^{-1})}.$$

where $j \in \mathbb{Z}$.

By Lemma 3.1 and a direct computation, we can get

$$(3.25) \quad q^{-j}e(j-2) \cdot q^{-j}g(j) = q^2z^2 + c_1z + c_2 \quad \text{for large even } j,$$

where c_1 and c_2 are complex numbers, which are independent of j .

Using (3.21)—(3.24), we have

$$(3.26) \quad q^{-j}e(j-2) = -q^2 \frac{R_1}{R_2}, \quad q^{-j}g(j) = -\frac{T_1}{T_2}$$

where

$$\begin{aligned}
R_1 &:= (z + [\alpha - 2]q^{\alpha-2} + [3][\beta]q^{-2})(z + [\alpha + 1]q^{\alpha-3} - [2]q^{-2}) \times \\
&\quad \times (z + [\alpha + 2]q^{\alpha-2} - [2]q^{-2}) \\
&\quad - q^{2\alpha-9}[\alpha + 1][\alpha + 2]f(0), \\
R_2 &:= (z + [\alpha + 1]q^{\alpha-3} - [2]q^{-2})(z + [\alpha + 2]q^{\alpha-2} - [2]q^{-2}), \\
T_1 &:= (z + [\alpha]q^{\alpha} - [\beta]q^{-2})(z + [\alpha - 1]q^{\alpha-1} + [2][\beta]q^{-1}) \times \\
&\quad \times (z + [\alpha - 2]q^{\alpha-2} + [2][\beta]q^{-3}) \\
&\quad - q^{-3}([\alpha - 1]q^{\alpha} + [2][\beta])([\alpha - 2]q^{\alpha} + [2][\beta]q^{-1})h(0), \\
T_2 &:= (z + [\alpha - 1]q^{\alpha-1} + [2][\beta]q^{-1})(z + [\alpha - 2]q^{\alpha-2} + [2][\beta]q^{-3}).
\end{aligned}$$

(3.25) implies that as the polynomials with respect to z , we have

$$(3.27) \quad R_2 T_2 \text{ divides } R_1 T_1$$

Now we have two cases to discuss:

• Case 1. $f(0)g(0) = 0$, in which case, either $f(0) = 0$ or $g(0) = 0$. If $f(0) = 0$, then (3.27) becomes

$$T_2 \text{ divides } (z + [\alpha - 2]q^{\alpha-2} + [3][\beta]q^{-2})T_1.$$

It follows that

$$(3.28) \quad - \frac{q^3([\alpha - 1]q^{\alpha} + [2][\beta])([\alpha - 2]q^{\alpha} + [2][\beta]q^{-1})h(0)}{T_2}$$

is a polynomial of z , hence, it is zero. Since $d_1(v_{-1}) \neq 0$ and $d_1(v_{-2}) \neq 0$, the coefficient of $h(0)$ in (3.28) is not zero. So we have to have $h(0) = 0$.

Similarly, if $h(0) = 0$, then we also have $f(0) = 0$.

Therefore, $f(0)g(0) = 0$ implies that $f(0) = g(0) = 0$. By (3.17)–(3.20), (3.16) is also true for $n = \pm 2$ and $j \in \mathbb{Z}$. This proves that $V = A(\alpha, \beta)$ because $U(W_q)$ is generated by $\{J^{\pm 1}, d_0, d_{\pm 1}, d_{\pm 2}\}$.

• Case 2. $f(0)g(0) \neq 0$. Since $d_{\pm 1}(v_j) \neq 0$ for all $j \in \mathbb{Z}$, the coefficients of $f(0)$ and $g(0)$ in R_1 and T_1 are non-zero. It follows from (3.27) that R_2 divides T_1 and T_2 divides R_1 , i.e.

$$T_1 = (z + [\alpha + 1]q^{\alpha-3} - [2]q^{-2})(z + [\alpha + 2]q^{\alpha-2} - [2]q^{-2})(z + G)$$

$$R_1 = (z + [\alpha - 1]q^{\alpha-1} + [2][\beta]q^{-1})(z + [\alpha - 2]q^{\alpha-2} + [2][\beta]q^{-3})(z + H)$$

where $G, H \in \mathbb{C}$. Comparing the coefficients of z^2 , we get

$$G = [\alpha - 2]q^{\alpha-2} + [3][\beta]q^{-2}, \quad H = q^\alpha[\alpha] - [\beta]q^{-2}.$$

Going back to (3.26), we have

$$\begin{aligned} & q^{-j}e(j) \\ &= -\frac{(z + [\alpha + 2]q^{\alpha+2} - [\beta]q^2)(z + [\alpha + 1]q^{\alpha+1} + [2][\beta]q^3)}{(z + [\alpha + 1]q^{\alpha+1})(z + [\alpha + 2]q^{\alpha+2})} \times \\ (3.29) \quad & \times (z + [\alpha]q^\alpha + [2][\beta]q), \end{aligned}$$

$$\begin{aligned} & q^{-j}g(j) \\ &= -\frac{(z + [\alpha + 1]q^{\alpha-3} - [2]q^{-2})(z + [\alpha + 2]q^{\alpha-2} - [2]q^{-2})}{(z + [\alpha - 1]q^{\alpha-1} + [2][\beta]q^{-1})(z + [\alpha - 2]q^{\alpha-2} + [2][\beta]q^{-3})} \times \\ (3.30) \quad & \times (z + [\alpha - 2]q^{\alpha-2} + [3][\beta]q^{-2}) \end{aligned}$$

for large even j . In particular, the rational function $q^{-j}e(j)$ of z and the rational function of the right side of (3.29) take the same values at infinitely many different points:

$$\{q^{-j}[j] \mid \text{for large even } j\}$$

It follows that (3.29) is true for all $j \in \mathbb{Z}$.

Similarly, (3.30) is also true for all $j \in \mathbb{Z}$.

Now we choose $a_0 = 1$ and $a_j \in \mathbb{C}^*$ for $j \in \mathbb{Z}$ such that

$$(3.31) \quad \frac{a_{j+k}}{a_{j+k+1}} = \frac{z + [\alpha + k + 1]q^{\alpha+k+1}}{z + [\alpha + k]q^{\alpha+k} + [2][\beta]q^{1+2k}} \quad \text{for } j, k \in \mathbb{Z}.$$

Set $u_j := a_j v_j$, we get

$$\begin{aligned} (3.32) \quad d_n(u_j) &= -q^j(z + [\alpha + n]q^{\alpha+n} + [1 - n][\beta]q^n)u_{n+j} \\ &= -([\alpha + n + j]q^{\alpha+n} + [1 - n][\beta]q^{n+j})u_{n+j} \end{aligned}$$

for $n = 0, \pm 1, \pm 2$ and $j \in \mathbb{Z}$.

For example, let us check that (3.32) is true for $n = 2$ and all $j \in \mathbb{Z}$. By (3.31), we have

$$(3.33) \quad \frac{a_j}{a_{j+2}} = \frac{(z + [\alpha + 1]q^{\alpha+1})(z + [\alpha + 2]q^{\alpha+2})}{(z + [\alpha]q^\alpha + [2][\beta]q)(z + [\alpha + 1]q^{\alpha+1} + [2][\beta]q^3)}$$

(3.29) and (3.33) imply that

$$\begin{aligned}
d_2(u_j) &= a_j(v_j) = a_j e(j) v_{j+2} = q^j \frac{a_j}{a_{j+2}} q^{-j} e(j) u_{j+2} \\
&= -q^j \frac{(z + [\alpha + 1]q^{\alpha+1})(z + [\alpha + 2]q^{\alpha+2})}{(z + [\alpha]q^\alpha + [2][\beta]q)(z + [\alpha + 1]q^{\alpha+1} + [2][\beta]q^3)} \times \\
&\quad \times \frac{(z + [\alpha + 2]q^{\alpha+2} - [\beta]q^2)(z + [\alpha + 1]q^{\alpha+1} + [2][\beta]q^3)}{(z + [\alpha + 1]q^{\alpha+1})(z + [\alpha + 2]q^{\alpha+2})} \times \\
&\quad \times (z + [\alpha]q^\alpha + [2][\beta]q) u_{j+2} \\
&= -q^j (z + [\alpha + 2]q^{\alpha+2} - [\beta]q^2) u_{j+2}.
\end{aligned}$$

Therefore, $V = \oplus_{n \in \mathbf{Z}} \mathbf{C} u_n = B(\alpha, \beta)$ by (3.32). ■

REFERENCES

1. B.L.Feigin and D.B.Fuchs, *Invariant skew-symmetric differential operators on the line and Verma modules over the Virasoro algebra*, Funct. Anal.Appl. (English translation) **16** (1982), 114-126.
2. V.G.Kac and A.K.Raina, "Highest weight representations of infinite dimensional Lie algebras," World Scientific Publishing Co Pte Ltd, 1987.
3. I.Kaplansky, *The Virasoro algebra*, Commun.Math.Phys. **86** (1982), 49-54.
4. I.Kaplansky and L.J.Santharoubane, *Harish-Chandra modules over the Virasoro algebra*, Proc.Conf. "Infinite dimensional Lie groups" Publications MSRI.
5. R.V.Moody and A.Pianzola, "Lie algebras with triangular decomposition," preprint.

CONCLUSIONS AND SOME OPEN PROBLEMS

After finishing this thesis, I become aware that a group of physicists have also begun to study q -deformations of the Virasoro algebra in their papers: [1], [3], [4], [5], [6] and [9]. Comparing these physicists' papers with my thesis, I find the following facts:

1. In spite of physicists have several methods of constructing the quantum Witt algebra, none of these methods is the same as my method of constructing the quantum Witt algebra.

2. Although the authors of [1], of [4] and I independently discovered the q -deformation of the Virasoro algebra under the condition that q is not a root of unity, the methods of computing the q -analogue of the Virasoro algebra central term are the same one, which is a q -version of the method of computing the Virasoro algebra central term in [2].

3. Physicists' work in [1], [3], [4], [5], [6] and [9] only overlap some work in Chapter 1 of this thesis.

4. Let $U_q(W)$ be the associative algebra generated by $\{d_m \mid m \in \mathbb{Z}\}$ with relations:

$$q^{m-n}d_md_n - q^{n-m}d_nd_m = [m-n]d_{m+n} \quad \text{for } m, n \in \mathbb{Z}.$$

This algebra was constructed by T. L. Curtright and C. K. Zachos in [6]. Using Definition 1.2 in Chapter 3 to define a $\mathbb{Z}$ -graded $U_q(W)$ -module, I note that $V = \bigoplus_{n \in \mathbb{Z}} V_n$ is a $\mathbb{Z}$ -graded $U_q(W)$ -module if and only if V is a $\mathbb{Z}$ -graded $U(W_q)$ -module with $J \mid V_n = q^n \cdot id$. So we can restate those results in Chapter 3 in terms of the algebra $U_q(W)$.

The facts above show that the work in this thesis has physics background. Recently, I have found that quantizing the Kac Conjecture proved by C. Martin and A. Piard in [8] is in connection with q -Gamma functions. Therefore, studying the quantization of the representation theory of the Virasoro algebra is closely related to both physics and some other areas in mathematics. The work in Chapter 3 is just a beginning on quantizing the representation theory of the Virasoro algebra, but it makes me believe that the work produced in developing the q -analogue of the representation theory of the Virasoro algebra will interest both physicists and mathematicians. In addition to this, studying the structure of quantum Lie algebras is also a quite interesting problem. I would like to conclude my thesis by proposing the following open problems about the structure of quantum Lie algebras:

Problem 1: Classify all complex finite dimensional $\mathbb{Z}$ -graded simple quantum Lie algebras.

Problem 2: Let $L = \bigoplus_{n \in \mathbb{Z}} L_n$ be a quantum Lie algebra. If $q^4 \neq 1$, then L_{-1} is a module over the Lie algebra L_0 . Therefore, one natural question is that finding the analogue of the Theorem in [7] for quantum Lie algebras. In other words, find all complex infinite dimensional $\mathbb{Z}$ -graded simple quantum Lie algebras $L = \bigoplus_{n \in \mathbb{Z}} L_n$ which satisfy the following conditions:

- (1) $\dim(L_n) < \infty$, for all $n \in \mathbb{Z}$.
- (2) $\lim_{n \rightarrow \infty} \frac{\ln(\dim(L_n))}{\ln |n|} < \infty$.
- (3) $L_{-1} \oplus L_0 \oplus L_1$ generates L and the module L_{-1} over Lie algebra L_0 is irreducible.

Problem 3: Is the quantum Witt algebra the unique complex infinite dimensional graded simple quantum Lie algebra $L = \bigoplus_{n \in \mathbb{Z}} L_n$ satisfying the conditions (1) and (2) above if $q^r \neq 1$ for $r=3,4,6,8$ and $L_{-1} \oplus L_0 \oplus L_1$ does not generate L ?

REFERENCES

1. N.Aizawa and H.Sato, *q-deformation of the Virasoro algebra with central extension*, Physics letters **256**, no.2 (1991), 185-190.
2. G.M.Benkart and R.V.Moody, *Derivations, central extensions and affine Lie algebras*, Algebras, Groups and Geometries **3** (1986), 456-492.
3. M.Chaichian, D.Ellinas and Z.Popowicz, *Quantum conformal algebra with central extension*, Physics Letters **248**, no.1,2 (1990), 95-99.
4. M.Chaichian, A.P.Isaev, J.Lukierski, Z.Popowicz and P.Presnajder, *q-deformations of Virasoro algebra and conformal dimensions*, Physics Letters **262**, no.1 (1991), 32-38.
5. M.Chaichian, P.Kulish and J.Lukierski, *q-deformed Jacobi identity, q-Oscillators and q-deformed infinite-dimensional algebras*, Physics Letters **237**, no.3,4 (1990), 401-406.
6. T.L.Curtright and C.K.Zachos, *Deforming maps for quantum algebras*, Physics Letters **243**, no.3 (1990), 237-244.
7. V.G.Kac, *Highest weight representations of infinite dimensional Lie algebras*, Proc. ICM. Helsinki (1978).
8. C.Martin and A.Piard, *Indecomposable modules over the Virasoro Lie algebra and a conjecture of V.Kac*, Commu. Math. Phys. **137** (1991), 109-132.
9. A.P.Polychronakos, *Consistency conditions and representations of a q-deformed Virasoro algebra*, Physics Letters **256**, no.1 (1991), 35-40.